

01/21/00  
3c759 U.S. PTO

01-24-00

A  
Box  
Seq

<b>UTILITY PATENT APPLICATION TRANSMITTAL</b>  <i>(Only for new nonprovisional applications under 37 CFR 1.53(b))</i>	Attorney Docket No.	L0461/7073 (JRV)
	First Named Inventor or Application Identifier	
	Ali Gure et al.	
	Express Mail Label No.	EL024661768US
	Date of Deposit	January 21, 2000

<b>APPLICATION ELEMENTS</b> <i>See MPEP chapter 600 concerning utility patent application contents</i>	<b>ADDRESS</b> <b>TO:</b> Box Patent Application Assistant Commissioner for Patents Washington, DC 20231
1. <input type="checkbox"/> Fee Transmittal Form <i>(Submit an original, and a duplicate for fee processing)</i>  2. <input checked="" type="checkbox"/> <b>Application</b> [Total pages 81] 64 - pages specification 1 - page abstract 16 - pages claims      116 - Total claims  3. <input checked="" type="checkbox"/> <b>Drawing(s)</b> (35 USC 113) [Total sheets 1] <input type="checkbox"/> Informal <input checked="" type="checkbox"/> Formal    [Total drawings 1]  4. <input type="checkbox"/> Oath or Declaration      [Total pages ] a. <input type="checkbox"/> Newly executed (original or copy) b. <input type="checkbox"/> Copy from a prior application (37 CFR 1.63(d)) <i>(for continuation/divisional with Box 17 completed)</i> [Note Box 5 below] i. <input type="checkbox"/> <b>DELETION OF INVENTOR(S)</b> Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b).  5. <input type="checkbox"/> Incorporation by Reference <i>(usable if Box 4b is checked)</i> The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.	6. <input type="checkbox"/> Microfiche Computer Program (Appendix)  7. <input checked="" type="checkbox"/> <b>Nucleotide and/or Amino Acid Sequence Submission</b> (if applicable, all necessary) a. <input checked="" type="checkbox"/> Computer Readable Copy b. <input checked="" type="checkbox"/> Paper Copy (identical to computer copy) c. <input checked="" type="checkbox"/> Statement verifying identity of above copies  <b>ACCOMPANYING APPLICATION PARTS</b>  8. <input type="checkbox"/> Assignment Papers/cover sheet & documents(s) 9. <input type="checkbox"/> 37 CFR 3.73(b) Statement <i>(when there is an assignee)</i> <input type="checkbox"/> Power of Attorney 10. <input type="checkbox"/> English Translation of Document <i>(if applicable)</i> 11. <input type="checkbox"/> Information Disclosure Statement PTO-1449 <input type="checkbox"/> Copies of IDS Citations 12. <input type="checkbox"/> Preliminary Amendment 13. <input checked="" type="checkbox"/> <b>Return Receipt Postcard</b> (MPEP 503) <i>(Should be specifically itemized)</i> 14. <input type="checkbox"/> Small Entity Statement(s) <input type="checkbox"/> Statement filed in prior application, Status still proper and desired 15. <input type="checkbox"/> Certified Copy of Priority Document(s) <i>(if foreign priority is claimed)</i>
16. Other:	
17. If a <b>CONTINUING APPLICATION</b> , check appropriate box and supply the requisite information:  <input type="checkbox"/> Continuation <input type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior application No.:  <input type="checkbox"/> Cancel in this application original claims of the prior application before calculating the filing fee.  <input type="checkbox"/> Amend the specification by inserting before the first line the sentence:  This application is a <input type="checkbox"/> continuation <input type="checkbox"/> divisional of application serial no. , filed , entitled , and now .	

18. CORRESPONDENCE ADDRESS					
Correspondence address below					
ATTORNEY'S NAME	John R. Van Amsterdam, Reg. No. 40,212				
NAME	Wolf, Greenfield & Sacks, P.C.				
ADDRESS	600 Atlantic Avenue				
CITY	Boston	STATE	MA	ZIP	02210
COUNTRY	USA	TELEPHONE	(617) 720-3500	FAX	(617) 720-2441

19. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED	
NAME	John R. Van Amsterdam, Reg. No. 40,212
SIGNATURE	<i>John R. Van Amsterdam</i>
DATE	<i>JANUARY 21, 2000</i>

**SMALL CELL LUNG CANCER ASSOCIATED ANTIGENS AND USES THEREFOR**

**Field of the Invention**

The invention relates to nucleic acids and encoded polypeptides which are cancer  
5 associated antigens expressed in patients afflicted with a variety of cancers. The invention  
also relates to agents which bind the nucleic acids or polypeptides. The nucleic acid  
molecules, polypeptides coded for by such molecules and peptides derived therefrom, as well  
as related antibodies and cytolytic T lymphocytes, are useful, *inter alia*, in diagnostic and  
therapeutic contexts.

**Background of the Invention**

The mechanism by which T cells recognize foreign materials has been implicated in  
cancer. A number of cytolytic T lymphocyte (CTL) clones directed against autologous  
melanoma antigens, testicular antigens, and melanocyte differentiation antigens have been  
15 described. In many instances, the antigens recognized by these clones have been  
characterized.

The use of autologous CTLs for identifying tumor antigens requires that the target  
cells which express the antigens can be cultured *in vitro* and that stable lines of autologous  
CTL clones which recognize the antigen-expressing cells can be isolated and propagated.  
20 While this approach has worked well for melanoma antigens, other tumor types, such as  
epithelial cancers including breast and colon cancer, have proved refractory to the approach.

More recently another approach to the problem has been described by Sahin et al.  
(*Proc. Natl. Acad. Sci. USA* 92:11810-11813, 1995). According to this approach, autologous  
antisera are used to identify immunogenic protein antigens expressed in cancer cells by  
25 screening expression libraries constructed from tumor cell cDNA. Antigen-encoding clones  
so identified have been found to have elicited an high-titer humoral immune response in the  
patients from which the antisera were obtained. Such a high-titer IgG response implies helper  
T cell recognition of the detected antigen. These tumor antigens can then be screened for the  
presence of MHC/HLA class I and class II motifs and reactivity with CTLs.

30 Presently there is a need for additional cancer antigens for development of therapeutics  
and diagnosis applicable to a greater number of cancer patients having various cancers.

### **Summary of the Invention**

Autologous antibody screening has now been applied to small cell lung cancer using antisera from cancer patients. Numerous cancer associated antigens have been identified.

The invention provides, *inter alia*, isolated nucleic acid molecules, expression vectors  
5 containing those molecules and host cells transfected with those molecules. The invention also provides isolated proteins and peptides, antibodies to those proteins and peptides and CTLs which recognize the proteins and peptides. Fragments including functional fragments and variants of the foregoing also are provided. Kits containing the foregoing molecules additionally are provided. The foregoing can be used in the diagnosis, monitoring, research,  
10 or treatment of conditions characterized by the expression of one or more cancer associated antigens.

Prior to the present invention, only a handful of small cell lung cancer associated genes had been identified in the past 20 years. The invention involves the surprising  
15 discovery of several genes, some previously known and some previously unknown, which are expressed in individuals who have cancer. These individuals all have serum antibodies against the proteins (or fragments thereof) encoded by these genes. Thus, abnormally expressed genes are recognized by the host's immune system and therefore can form a basis for diagnosis, monitoring and therapy.

The invention involves the use of a single material, a plurality of different materials  
20 and even large panels and combinations of materials. For example, a single gene, a single protein encoded by a gene, a single functional fragment thereof, a single antibody thereto, etc. can be used in methods and products of the invention. Likewise, pairs, groups and even panels of these materials and optionally other cancer associated antigen genes and/or gene products can be used for diagnosis, monitoring and therapy. The pairs, groups or panels can  
25 involve 2, 3, 4, 5 or more genes, gene products, fragments thereof or agents that recognize such materials. A plurality of such materials are not only useful in monitoring, typing, characterizing and diagnosing cells abnormally expressing such genes, but a plurality of such materials can be used therapeutically. An example of the use of a plurality of such materials for the prevention, delay of onset, amelioration, etc. of cancer cells, which express or will  
30 express such genes prophylactically or acutely. Any and all combinations of the genes, gene products, and materials which recognize the genes and gene products can be tested and identified for use according to the invention. It would be far too lengthy to recite all such combinations; those skilled in the art, particularly in view of the teaching contained herein,

will readily be able to determine which combinations are most appropriate for which circumstances.

As will be clear from the following discussion, the invention has *in vivo* and *in vitro* uses, including for therapeutic, diagnostic, monitoring and research purposes. One aspect of the invention is the ability to fingerprint a cell expressing a number of the genes identified according to the invention by, for example, quantifying the expression of such gene products. Such fingerprints will be characteristic, for example, of the stage of the cancer, the type of the cancer, or even the effect in animal models of a therapy on a cancer. Cells also can be screened to determine whether such cells abnormally express the genes identified according to the invention.

The invention, in one aspect, is a method of diagnosing a disorder characterized by expression of a cancer associated antigen precursor coded for by a nucleic acid molecule. The method involves the steps of contacting a biological sample isolated from a subject with an agent that specifically binds to the nucleic acid molecule, an expression product thereof, or a fragment of an expression product thereof complexed with an MHC, preferably an HLA, molecule, wherein the nucleic acid molecule is a NA Group 1 nucleic acid molecule, and determining the interaction between the agent and the nucleic acid molecule, the expression product or fragment of the expression product as a determination of the disorder.

In one embodiment the agent is selected from the group consisting of (a) a nucleic acid molecule comprising NA Group 1 nucleic acid molecules or a fragment thereof, (b) a nucleic acid molecule comprising NA Group 3 nucleic acid molecules or a fragment thereof, (c) a nucleic acid molecule comprising NA Group 5 nucleic acid molecules or a fragment thereof, (d) an antibody that binds to an expression product, or a fragment thereof, of NA group 1 nucleic acids, (e) an antibody that binds to an expression product, or a fragment thereof, of NA group 3 nucleic acids, (f) an antibody that binds to an expression product, or a fragment thereof, of NA group 5 nucleic acids, (g) an agent that binds to a complex of an MHC, preferably HLA, molecule and a fragment of an expression product of a NA Group 1 nucleic acid, (h) an agent that binds to a complex of an MHC, preferably HLA, molecule and a fragment of an expression product of a NA group 3 nucleic acid, and (i) an agent that binds to a complex of an MHC, preferably HLA, molecule and a fragment of an expression product of a NA Group 5 nucleic acid.

The disorder may be characterized by expression of a plurality of cancer associated antigen precursors. Thus the methods of diagnosis may include use of a plurality of agents,

each of which is specific for a different human cancer associated antigen precursor (including at least one of the cancer associated antigen precursors disclosed herein), and wherein said plurality of agents is at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9 or at least 10 such agents.

5           In each of the above embodiments the disorder preferably is selected from the group consisting of lung cancers including small cell lung cancer and non-small cell lung cancer, melanoma, colon cancer, breast cancer, head and neck cancer, transitional cancer, leiomyosarcoma and synovial sarcoma.

10           In some embodiments, the nucleic acid molecule is selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids. Preferably the nucleic acid molecule is selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12.

15           In certain embodiments, the biological sample is isolated from a tissue selected from the group consisting of non-brain, non-testis, non-prostate, non-small intestine and non-colon tissues.

20           In another aspect the invention is a method for determining regression, progression or onset of a condition characterized by expression of abnormal levels of a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule. The method involves the steps of monitoring a sample, from a subject who has or is suspected of having the condition, for a parameter selected from the group consisting of (i) the protein, (ii) a peptide derived from the protein, (iii) an antibody which selectively binds the protein or peptide, and (iv) cytolytic T cells specific for a complex of the peptide derived from the protein and an MHC molecule, as a determination of regression, progression or onset of said condition. In one embodiment the sample is a body fluid, a body effusion or a tissue.

25           In another embodiment the step of monitoring comprises contacting the sample with a detectable agent selected from the group consisting of (a) an antibody which selectively binds the protein of (i), or the peptide of (ii), (b) a protein or peptide which binds the antibody of (iii), and (c) a cell which presents the complex of the peptide and MHC molecule of (iv). In a preferred embodiment the antibody, the protein, the peptide or the cell is labeled with a radioactive label or an enzyme. The sample in a preferred embodiment is assayed for the peptide.

          According to another embodiment the nucleic acid molecule is one of the following: a

NA Group 3 molecule or a NA Group 5 molecule. In still another embodiment, the nucleic acid molecule is selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids. Preferably the nucleic acid molecule is selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12.

In yet another embodiment the protein is a plurality of proteins, the parameter is a plurality of parameters, each of the plurality of parameters being specific for a different of the plurality of proteins, at least one of which is a cancer associated protein encoded by a NA group 1 molecule. In certain embodiments the protein is a plurality of proteins, at least one of which is encoded by SOX2 (SEQ ID NO:3) or ZIC2 (SEQ ID NO:5), and wherein the parameter is a plurality of parameters, each of the plurality of parameters being specific for a different of the plurality of proteins.

The invention in another aspect is a pharmaceutical preparation for a human subject. The pharmaceutical preparation includes an agent which when administered to the subject enriches selectively the presence of complexes of an HLA molecule and a human cancer associated antigen, and a pharmaceutically acceptable carrier, wherein the human cancer associated antigen is a fragment of a human cancer associated antigen precursor encoded by a nucleic acid molecule which comprises a NA Group 1 molecule. In one embodiment the nucleic acid molecule is a NA Group 3 nucleic acid molecule or a NA group 5 nucleic acid molecule.

The agent in one embodiment comprises a plurality of agents, each of which enriches selectively in the subject complexes of an HLA molecule and a different human cancer associated antigen. Preferably the plurality is at least two, at least three, at least four or at least 5 different such agents.

In certain embodiments, the agent comprises a plurality of agents, at least one of which is a nucleic acid molecule selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids, and preferably at least one of which is a nucleic acid molecule selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12, or an expression product thereof, and each of which enriches selectively in the subject complexes of an HLA molecule and a different human cancer associated antigen.

In another embodiment the agent is selected from the group consisting of (1) an isolated polypeptide comprising the human cancer associated antigen, or a functional variant

thereof, (2) an isolated nucleic acid operably linked to a promoter for expressing the isolated polypeptide, or functional variant thereof, (3) a host cell expressing the isolated polypeptide, or functional variant thereof, and (4) isolated complexes of the polypeptide, or functional variants thereof, and an HLA molecule.

5           The agent may be a cell expressing an isolated polypeptide. In one embodiment the agent is a cell expressing an isolated polypeptide comprising the human cancer associated antigen or a functional variant thereof. In another embodiment the agent is a cell expressing an isolated polypeptide comprising the human cancer associated antigen or a functional variant thereof, and wherein the cell expresses an HLA molecule that binds the polypeptide.

10       The cell can express one or both of the polypeptide and HLA molecule recombinantly. In preferred embodiments the cell is nonproliferative. In other preferred embodiments, the isolated polypeptide is or includes a polypeptide encoded by a nucleic acid molecule selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids, and preferably at least one of which is a  
15       nucleic acid molecule selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12. In yet another embodiment the agent is at least two, at least three, at least four or at least five different polypeptides, each representing a different human cancer associated antigen or functional variant thereof.

          The agent in one embodiment is a PP Group 2 polypeptide. In other embodiments the  
20       agent is a PP Group 3 polypeptide or a PP Group 4 polypeptide.

          In an embodiment each of the pharmaceutical preparations described herein also includes an adjuvant.

          According to another aspect the invention, a composition is provided which includes an isolated agent that binds selectively a PP Group 1 polypeptide. In separate embodiments  
25       the agent binds selectively to a polypeptide selected from the following: a PP Group 2 polypeptide, a PP Group 3 polypeptide, a PP Group 4 polypeptide, and a PP Group 5 polypeptide. In other embodiments, the agent is a plurality of different agents that bind selectively at least two, at least three, at least four, or at least five different such polypeptides.

          In each of the above described embodiments the agent may be an antibody. In a preferred  
30       embodiment, at least one of polypeptides is encoded by a nucleic acid molecule selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids, and preferably at least one of which is a nucleic acid molecule selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5,

SEQ ID NO:11 and SEQ ID NO:12, or a fragment thereof.

In another aspect the invention is a composition of matter composed of a conjugate of the agent of the above-described compositions of the invention and a therapeutic or diagnostic agent. Preferably the conjugate is of the agent and a therapeutic or diagnostic that is a toxin, particularly an antineoplastic.

The invention in another aspect is a pharmaceutical composition which includes an isolated nucleic acid molecule selected from the group consisting of: (1) NA Group 1 molecules, and (2) NA Group 2 molecules, and a pharmaceutically acceptable carrier. In one embodiment the isolated nucleic acid molecule comprises a NA Group 3 or NA Group 4 molecule. In another embodiment the isolated nucleic acid molecule comprises at least two isolated nucleic acid molecules coding for two different polypeptides, each polypeptide comprising a different cancer associated antigen. In preferred embodiments, at least one of the polypeptides is encoded by a nucleic acid molecule selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids, and preferably at least one of which is a nucleic acid molecule selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12.

Preferably the pharmaceutical composition also includes an expression vector with a promoter operably linked to the isolated nucleic acid molecule. In another embodiment the pharmaceutical composition also includes a host cell recombinantly expressing the isolated nucleic acid molecule.

According to another aspect of the invention a pharmaceutical composition is provided. The pharmaceutical composition includes an isolated polypeptide comprising a PP Group 1 or a PP Group 2 polypeptide, and a pharmaceutically acceptable carrier. In one embodiment the isolated polypeptide comprises a PP Group 3 or a PP Group 4 polypeptide.

In another embodiment the isolated polypeptide comprises at least two different polypeptides, each comprising a different cancer associated antigen at least one of which is encoded by a NA group 1 molecule as disclosed herein. In certain embodiments at least one of the polypeptides is encoded by a nucleic acid molecule selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids, and preferably at least one of which is a nucleic acid molecule selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12. In separate embodiments the isolated polypeptides are selected from the

following: PP Group 3 polypeptides or HLA binding fragments thereof and PP Group 5 polypeptides or HLA binding fragments thereof.

In an embodiment each of the pharmaceutical compositions described herein also includes an adjuvant.

5 Another aspect the invention is an isolated nucleic acid molecule comprising a NA Group 3 molecule. Another aspect the invention is an isolated nucleic acid molecule comprising a NA Group 4 molecule.

The invention in another aspect is an isolated nucleic acid molecule selected from the group consisting of (a) a fragment of a nucleic acid selected from the group of nucleic acid  
10 molecules consisting of SEQ ID Nos numbered below and comprising all nucleic acid sequences among SEQ ID Nos:3-17, of sufficient length to represent a sequence unique within the human genome, and identifying a nucleic acid encoding a human cancer associated antigen precursor, (b) complements of (a), provided that the fragment includes a sequence of contiguous nucleotides which is not identical to any sequence selected from the sequence  
15 group consisting of (1) sequences having the GenBank accession numbers of Table 4, (2) complements of (1), and (3) fragments of (1) and (2).

In one embodiment the sequence of contiguous nucleotides is selected from the group consisting of: (1) at least two contiguous nucleotides nonidentical to the sequences in Table 4, (2) at least three contiguous nucleotides nonidentical to the sequences in Table 4, (3) at least  
20 four contiguous nucleotides nonidentical to the sequences in Table 4, (4) at least five contiguous nucleotides nonidentical to the sequences in Table 4, (5) at least six contiguous nucleotides nonidentical to the sequences in Table 4, or (6) at least seven contiguous nucleotides nonidentical to the sequences in Table 4.

In another embodiment the fragment has a size selected from the group consisting of at  
25 least: 8 nucleotides, 10 nucleotides, 12 nucleotides, 14 nucleotides, 16 nucleotides, 18 nucleotides, 20, nucleotides, 22 nucleotides, 24 nucleotides, 26 nucleotides, 28 nucleotides, 30 nucleotides, 50 nucleotides, 75 nucleotides, 100 nucleotides, 200 nucleotides, 1000 nucleotides and every integer length therebetween.

In yet another embodiment the molecule encodes a polypeptide which, or a fragment  
30 of which, binds a human HLA receptor or a human antibody.

Another aspect of the invention is an expression vector comprising an isolated nucleic acid molecule of the invention described above operably linked to a promoter.

According to one aspect the invention is an expression vector comprising a nucleic

acid operably linked to a promoter, wherein the nucleic acid is a NA Group 1 or Group 2 molecule. In another aspect the invention is an expression vector comprising a NA Group 1 or Group 2 molecule and a nucleic acid encoding an MHC, preferably HLA, molecule.

5 In yet another aspect the invention is a host cell transformed or transfected with an expression vector of the invention described above.

In another aspect the invention is a host cell transformed or transfected with an expression vector comprising an isolated nucleic acid molecule of the invention described above operably linked to a promoter, or an expression vector comprising a nucleic acid operably linked to a promoter, wherein the nucleic acid is a NA Group 1 or 2 molecule and  
10 further comprising a nucleic acid encoding HLA.

According to another aspect of the invention an isolated polypeptide encoded by the isolated nucleic acid molecules the invention, described above, is provided. These include PP Group 1-5 polypeptides. The invention also includes a fragment of the polypeptide which is immunogenic. In one embodiment the fragment, or a portion of the fragment, binds HLA or a  
15 human antibody. In still another aspect the invention provides as isolated polypeptide comprising a fragment of a polypeptide selected from the group consisting of ZIC2, SOX1, SOX2, SOX3 and SOX21 polypeptides, which is immunogenic, wherein the polypeptide is not a full-length ZIC1, SOX1, SOX2, SOX3 or SOX21 polypeptide.

The invention includes in another aspect an isolated fragment of a human cancer  
20 associated antigen precursor which, or portion of which, binds HLA or a human antibody, wherein the precursor is encoded by a nucleic acid molecule that is a NA Group 1 molecule. In one embodiment the fragment is part of a complex with HLA. In another embodiment the fragment is between 8 and 12 amino acids in length. In another embodiment the invention includes an isolated polypeptide comprising a fragment of the polypeptide of sufficient length  
25 to represent a sequence unique within the human genome and identifying a polypeptide that is a human cancer associated antigen precursor.

According to another aspect of the invention a kit for detecting the presence of the expression of a cancer associated antigen precursor is provided. The kit includes a pair of isolated nucleic acid molecules each of which consists essentially of a molecule selected from  
30 the group consisting of (a) a 12-32 nucleotide contiguous segment of the nucleotide sequence of any of the NA Group 1 molecules and (b) complements of ("a"), wherein the contiguous segments are nonoverlapping. In one embodiment the pair of isolated nucleic acid molecules is constructed and arranged to selectively amplify an isolated nucleic acid molecule that is a

NA Group 3 molecule. Preferably, the pair amplifies a human NA Group 3 molecule.

According to another aspect of the invention a method for treating a subject with a disorder characterized by expression of a human cancer associated antigen precursor is provided. The method includes the step of administering to the subject an amount of an agent, which enriches selectively in the subject the presence of complexes of an HLA molecule and a human cancer associated antigen, effective to ameliorate the disorder, wherein the human cancer associated antigen is a fragment of a human cancer associated antigen precursor encoded by a nucleic acid molecule selected from the group consisting of (a) a nucleic acid molecule comprising NA group 1 nucleic acid molecules, (b) a nucleic acid molecule comprising NA group 3 nucleic acid molecules, (c) a nucleic acid molecule comprising NA group 5 nucleic acid molecules.

In one embodiment the disorder is characterized by expression of a plurality of human cancer associated antigen precursors and wherein the agent is a plurality of agents, each of which enriches selectively in the subject the presence of complexes of an HLA molecule and a different human cancer associated antigen. Preferably the plurality is at least 2, at least 3, at least 4, or at least 5 such agents. In a preferred embodiment, at least one of the human cancer associated antigens is a polypeptide encoded by a nucleic acid molecule selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids, and preferably at least one of which is a nucleic acid molecule selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12, or a fragment thereof.

In another embodiment the agent is an isolated polypeptide selected from the group consisting of PP Group 1, PP Group 2, PP Group 3, PP Group 4, and PP group 5 polypeptides.

In yet another embodiment the disorder is cancer.

According to another aspect the invention is a method for treating a subject having a condition characterized by expression of a cancer associated antigen precursor in cells of the subject. The method includes the steps of (i) removing an immunoreactive cell containing sample from the subject, (ii) contacting the immunoreactive cell containing sample to the host cell under conditions favoring production of cytolytic T cells against a human cancer associated antigen which is a fragment of the precursor, (iii) introducing the cytolytic T cells to the subject in an amount effective to lyse cells which express the human cancer associated antigen, wherein the host cell is transformed or transfected with an expression vector

comprising an isolated nucleic acid molecule operably linked to a promoter, the isolated nucleic acid molecule being selected from the group of nucleic acid molecules consisting of NA Group 1, NA Group 2, NA Group 3, NA Group 4, NA Group 5.

In one embodiment the host cell recombinantly expresses an HLA molecule which binds the human cancer associated antigen. In another embodiment the host cell endogenously expresses an HLA molecule which binds the human cancer associated antigen.

The invention includes in another aspect a method for treating a subject having a condition characterized by expression of a cancer associated antigen precursor in cells of the subject. The method includes the steps of (i) identifying a nucleic acid molecule expressed by the cells associated with said condition, wherein said nucleic acid molecule is a NA Group 1 molecule (ii) transfecting a host cell with a nucleic acid selected from the group consisting of (a) the nucleic acid molecule identified, (b) a fragment of the nucleic acid identified which includes a segment coding for a cancer associated antigen, (c) deletions, substitutions or additions to (a) or (b), and (d) degenerates of (a), (b), or (c); (iii) culturing said transfected host cells to express the transfected nucleic acid molecule, and; (iv) introducing an amount of said host cells or an extract thereof to the subject effective to increase an immune response against the cells of the subject associated with the condition. Preferably, the antigen is a human antigen and the subject is a human. In certain preferred embodiments the nucleic acid molecule is selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids, and preferably at least one of which is a nucleic acid molecule is selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12.

In one embodiment the method also includes the step of (a) identifying an MHC molecule which presents a portion of an expression product of the nucleic acid molecule, wherein the host cell expresses the same MHC molecule as identified in (a) and wherein the host cell presents an MHC binding portion of the expression product of the nucleic acid molecule.

In another embodiment the method also includes the step of treating the host cells to render them non-proliferative.

In yet another embodiment the immune response comprises a B-cell response or a T cell response. Preferably the response is a T-cell response which comprises generation of cytolytic T-cells specific for the host cells presenting the portion of the expression product of the nucleic acid molecule or cells of the subject expressing the human cancer associated

antigen.

In another embodiment the nucleic acid molecule is a NA Group 3 molecule.

Another aspect of the invention is a method for treating or diagnosing or monitoring a subject having a condition characterized by expression of an abnormal amount of a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule. The method includes the step of administering to the subject an antibody which specifically binds to the protein or a peptide derived therefrom, the antibody being coupled to a therapeutically useful agent, in an amount effective to treat the condition.

In one embodiment the antibody is a monoclonal antibody. Preferably the monoclonal antibody is a chimeric antibody or a humanized antibody.

In another aspect the invention is a method for treating a condition characterized by expression in a subject of abnormal amounts of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule. The method involves the step of administering to a subject at least one of the pharmaceutical compositions of the invention described above in an amount effective to prevent, delay the onset of, or inhibit the condition in the subject. In one embodiment the condition is cancer. In another embodiment the method includes the step of first identifying that the subject expresses in a tissue abnormal amounts of the protein.

The invention in another aspect is a method for treating a subject having a condition characterized by expression of abnormal amounts of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule. The method includes the steps of (i) identifying cells from the subject which express abnormal amounts of the protein; (ii) isolating a sample of the cells; (iii) cultivating the cells, and (iv) introducing the cells to the subject in an amount effective to provoke an immune response against the cells.

In one embodiment the method includes the step of rendering the cells non-proliferative, prior to introducing them to the subject.

In another aspect the invention is a method for treating a pathological cell condition characterized by abnormal expression of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule. The method includes the step of administering to a subject in need thereof an effective amount of an agent which inhibits the expression or activity of the protein.

In one embodiment the agent is an inhibiting antibody which selectively binds to the protein and wherein the antibody is a monoclonal antibody, a chimeric antibody, a humanized antibody or a fragment thereof. In another embodiment the agent is an antisense nucleic acid

molecule which selectively binds to the nucleic acid molecule which encodes the protein. In yet another important embodiment the nucleic acid molecule is a NA Group 3 nucleic acid molecule. In other preferred embodiments, the nucleic acid molecule is a nucleic acid molecule selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids, and preferably at least one of which is a nucleic acid molecule selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12.

The invention includes in another aspect a composition of matter useful in stimulating an immune response to a plurality of proteins encoded by nucleic acid molecules that are NA Group 1 molecules. The composition is a plurality of peptides derived from the amino acid sequences of the proteins, wherein the peptides bind to one or more MHC molecules presented on the surface of the cells which express an abnormal amount of the protein. In preferred embodiments, at least one of the proteins is encoded by a nucleic acid molecule selected from the group consisting of SOX2 nucleic acids, SOX1 nucleic acids, ZIC2 nucleic acids, SOX3 nucleic acids and SOX21 nucleic acids, and preferably at least one of which is a nucleic acid molecule selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:11 and SEQ ID NO:12.

In one embodiment at least a portion of the plurality of peptides bind to MHC molecules and elicit a cytolytic response thereto. In another embodiment the composition of matter includes an adjuvant. In another embodiment the adjuvant is a saponin, GM-CSF, or an interleukin. In still another embodiment, the compositions also includes at least one peptide useful in stimulating an immune response to at least one protein which is not encoded by nucleic acid molecules that are NA Group 1 molecules, wherein the at least one peptide binds to one or more MHC molecules.

According to another aspect the invention is an isolated antibody which selectively binds to a complex of: (i) a peptide derived from a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule and (ii) and an MHC molecule to which binds the peptide to form the complex, wherein the isolated antibody does not bind to (i) or (ii) alone.

In one embodiment the antibody is a monoclonal antibody, a chimeric antibody, a humanized antibody or a fragment thereof.

The invention also involves the use of the genes, gene products, fragments thereof, agents which bind thereto, and so on in the preparation of medicaments. A particular medicament is for treating cancer and a more particular medicament is for treating small cell

lung cancer.

For all of the foregoing, preferred disorders include cancers, particularly lung cancers including small cell lung cancer and non-small cell lung cancer, melanoma, colon cancer, breast cancer, head and neck cancer, transitional cancer, leiomyosarcoma and synovial sarcoma. Preferred tissues include non-brain, non-testis, non-prostate, non-small intestine and non-colon tissues.

These and other aspects of the invention will be described in further detail in connection with the detailed description of the invention.

#### **Brief Description of the Figure**

Fig. 1 shows the alignment of predicted protein sequences of SOX1, 2, 3 and 21 (GenBank accession numbers O00570, P48431, P41225, AAC95381.1, respectively; SEQ ID Nos:18-21). Sequences encoded within the SEREX-isolated clones are in bold face type, and sequences absent in these clones are in gray italics. The DNA-binding HMG domain is boxed. Amino acids identical between three and four SOX proteins are highlighted in two shades of gray.

#### **Detailed Description of the Invention**

In the above summary and in the ensuing description, lists of sequences are provided. The lists are meant to embrace each single sequence separately, two or more sequences together where they form a part of the same gene, any combination of two or more sequences which relate to different genes, including and up to the total number on the list, as if each and every combination were separately and specifically enumerated. Likewise, when mentioning fragment size, it is intended that a range embrace the smallest fragment mentioned to the full-length of the sequence (less one nucleotide or amino acid so that it is a fragment), each and every fragment length intended as if specifically enumerated. Thus, if a fragment could be between 10 and 15 in length, it is explicitly meant to mean 10, 11, 12, 13, 14, or 15 in length.

The summary and the claims mention antigen precursors and antigens. As used in the summary and in the claims, a precursor is substantially the full-length protein encoded by the coding region of the isolated DNA and the antigen is a peptide which complexes with MHC, preferably HLA, and which participates in the immune response as part of that complex. Such antigens are typically 9 amino acids long, although this may vary slightly.

As used herein, a subject is a human, non-human primate, cow, horse, pig, sheep, goat,

dog, cat or rodent. In all embodiments human cancer antigens and human subjects are preferred.

The present invention in one aspect involves the cloning of cDNAs encoding human small cell lung cancer associated antigen precursors using autologous antisera of subjects having cancer. The sequences of the clones representing genes identified according to the methods described herein are presented in the attached Sequence Listing. Of the foregoing, it can be seen that some of the clones are novel but may have some homology to sequences deposited in databases (mainly EST sequences). Nevertheless, the entire gene sequence was not previously known. In some cases no function was suspected and in other cases, even if a function was suspected, it was not known that the gene was associated with cancer. In all cases, it was not known or suspected that the gene encoded a cancer antigen which reacted with antibody from autologous sera. Analysis of the clone sequences by comparison to nucleic acid and protein databases determined that still other of the clones surprisingly are closely related to other previously-cloned genes. The sequences of these related genes is also presented in the Sequence Listing. The nature of the foregoing genes as encoding antigens recognized by the immune systems of cancer patients is, of course, unexpected.

The invention thus involves in one aspect cancer associated antigen polypeptides, genes encoding those polypeptides, functional modifications and variants of the foregoing, useful fragments of the foregoing, as well as diagnostics and therapeutics relating thereto.

Homologs and alleles of the cancer associated antigen nucleic acids of the invention can be identified by conventional techniques. Thus, an aspect of the invention is those nucleic acid sequences which code for cancer associated antigen precursors. Because this application contains so many sequences, the following chart is provided to identify the various groups of sequences discussed in the claims and in the summary:

#### Nucleic Acid Sequences

NA Group 1. (a) nucleic acid molecules which hybridize under stringent conditions to a molecule consisting of a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 3-17 and which code for a cancer associated antigen precursor,

(b) deletions, additions and substitutions which code for a respective cancer associated antigen precursor,

(c) nucleic acid molecules that differ from the nucleic acid molecules of (a) or (b) in codon sequence due to the degeneracy of the genetic code, and

(d) complements of (a), (b) or (c).

5 NA Group 2. Fragments of NA Group 1, which code for a polypeptide which, or a portion of which, binds a MHC molecule to form a complex recognized by an autologous antibody or lymphocyte.

NA Group 3. The subset of NA Group 1 where the nucleotide sequence is selected from the group consisting of:

10 (a) previously unknown human nucleic acids coding for a human cancer associated antigen precursor set forth as SEQ ID NO:17,

(b) deletions, additions and substitutions which code for a respective human cancer associated antigen precursor,

(c) nucleic acid molecules that differ from the nucleic acid molecules of (a) or (b) in codon sequence due to the degeneracy of the genetic code, and

15 (d) complements of (a), (b) or (c).

NA Group 4. Fragments of NA Group 3, which code for a polypeptide which, or a portion of which, binds to a MHC molecule to form a complex recognized by an autologous antibody or lymphocyte.

20 NA Group 5. A subset of NA Group 1, comprising human cancer associated antigens that react with allogeneic cancer antisera.

#### Polypeptide Sequences

25 PP Group 1. Polypeptides encoded by NA Group 1.

PP Group 2. Polypeptides encoded by NA Group 2

PP Group 3. Polypeptides encoded by NA Group 3.

PP Group 4. Polypeptides encoded by NA Group 4.

PP Group 5. Polypeptides encoded by NA Group 5.

30

The term “stringent conditions” as used herein refers to parameters with which the art is familiar. Nucleic acid hybridization parameters may be found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds.,

Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. More specifically, stringent conditions, as used herein, refers, for example, to hybridization at 65°C in hybridization buffer (3.5 x SSC, 0.02% Ficoll, 0.02% polyvinyl pyrrolidone, 0.02% Bovine Serum Albumin, 2.5mM NaH<sub>2</sub>PO<sub>4</sub>(pH7), 0.5% SDS, 2mM EDTA). SSC is 0.15M sodium chloride/0.15M sodium citrate, pH7; SDS is sodium dodecyl sulphate; and EDTA is ethylenediaminetetracetic acid. After hybridization, the membrane upon which the DNA is transferred is washed, for example, in 2 x SSC at room temperature and then at 0.1 - 0.5 x SSC/0.1 x SDS at temperatures up to 68°C.

There are other conditions, reagents, and so forth which can be used, which result in a similar degree of stringency. The skilled artisan will be familiar with such conditions, and thus they are not given here. It will be understood, however, that the skilled artisan will be able to manipulate the conditions in a manner to permit the clear identification of homologs and alleles of cancer associated antigen nucleic acids of the invention (e.g., by using lower stringency conditions). The skilled artisan also is familiar with the methodology for screening cells and libraries for expression of such molecules which then are routinely isolated, followed by isolation of the pertinent nucleic acid molecule and sequencing.

In general homologs and alleles typically will share at least 75% nucleotide identity and/or at least 90% amino acid identity to the sequences of cancer associated antigen nucleic acid and polypeptides, respectively, in some instances will share at least 90% nucleotide identity and/or at least 95% amino acid identity and in still other instances will share at least 95% nucleotide identity and/or at least 99% amino acid identity. The homology can be calculated using various, publicly available software tools developed by NCBI (Bethesda, Maryland) that can be obtained through the internet (<ftp://ncbi.nlm.nih.gov/pub/>). Exemplary tools include the BLAST system available at <http://www.ncbi.nlm.nih.gov>, using default settings. Pairwise and ClustalW alignments (BLOSUM30 matrix setting) as well as Kyte-Doolittle hydropathic analysis can be obtained using the MacVector sequence analysis software (Oxford Molecular Group). Watson-Crick complements of the foregoing nucleic acids also are embraced by the invention.

In screening for cancer associated antigen genes, a Southern blot may be performed using the foregoing conditions, together with a radioactive probe. After washing the membrane to which the DNA is finally transferred, the membrane can be placed against X-ray film to detect the radioactive signal. In screening for the expression of cancer associated

antigen nucleic acids, Northern blot hybridizations using the foregoing conditions (see also the Examples) can be performed on samples taken from breast cancer patients or subjects suspected of having a condition characterized by expression of breast cancer associated antigen genes. Amplification protocols such as polymerase chain reaction using primers which hybridize to the sequences presented also can be used for detection of the cancer associated antigen genes or expression thereof.

The small cell lung cancer associated genes correspond to SEQ ID NOs. 3-17. The preferred cancer associated antigens for the methods of diagnosis disclosed herein are those which were found to react with allogeneic cancer antisera (i.e. NA Group 5). Especially preferred are the *ZIC2* and *SOX Group B* sequences (SEQ ID Nos: 3, 4, 5, 11 and 12). Encoded polypeptides (e.g., SEQ ID NOS:18-22), peptides and antisera thereto are also preferred for diagnosis.

The invention also includes degenerate nucleic acids which include alternative codons to those present in the native materials. For example, serine residues are encoded by the codons TCA, AGT, TCC, TCG, TCT and AGC. Each of the six codons is equivalent for the purposes of encoding a serine residue. Thus, it will be apparent to one of ordinary skill in the art that any of the serine-encoding nucleotide triplets may be employed to direct the protein synthesis apparatus, *in vitro* or *in vivo*, to incorporate a serine residue into an elongating breast cancer associated antigen polypeptide. Similarly, nucleotide sequence triplets which encode other amino acid residues include, but are not limited to: CCA, CCC, CCG and CCT (proline codons); CGA, CGC, CGG, CGT, AGA and AGG (arginine codons); ACA, ACC, ACG and ACT (threonine codons); AAC and AAT (asparagine codons); and ATA, ATC and ATT (isoleucine codons). Other amino acid residues may be encoded similarly by multiple nucleotide sequences. Thus, the invention embraces degenerate nucleic acids that differ from the biologically isolated nucleic acids in codon sequence due to the degeneracy of the genetic code.

The invention also provides modified nucleic acid molecules which include additions, substitutions and deletions of one or more nucleotides. In preferred embodiments, these modified nucleic acid molecules and/or the polypeptides they encode retain at least one activity or function of the unmodified nucleic acid molecule and/or the polypeptides, such as antigenicity, enzymatic activity, receptor binding, formation of complexes by binding of peptides by MHC class I and class II molecules, etc. In certain embodiments, the modified nucleic acid molecules encode modified polypeptides, preferably polypeptides having

conservative amino acid substitutions as are described elsewhere herein. The modified nucleic acid molecules are structurally related to the unmodified nucleic acid molecules and in preferred embodiments are sufficiently structurally related to the unmodified nucleic acid molecules so that the modified and unmodified nucleic acid molecules hybridize under stringent conditions known to one of skill in the art.

For example, modified nucleic acid molecules which encode polypeptides having single amino acid changes can be prepared. Each of these nucleic acid molecules can have one, two or three nucleotide substitutions exclusive of nucleotide changes corresponding to the degeneracy of the genetic code as described herein. Likewise, modified nucleic acid molecules which encode polypeptides having two amino acid changes can be prepared which have, e.g., 2-6 nucleotide changes. Numerous modified nucleic acid molecules like these will be readily envisioned by one of skill in the art, including for example, substitutions of nucleotides in codons encoding amino acids 2 and 3, 2 and 4, 2 and 5, 2 and 6, and so on. In the foregoing example, each combination of two amino acids is included in the set of modified nucleic acid molecules, as well as all nucleotide substitutions which code for the amino acid substitutions. Additional nucleic acid molecules that encode polypeptides having additional substitutions (i.e., 3 or more), additions or deletions (e.g., by introduction of a stop codon or a splice site(s)) also can be prepared and are embraced by the invention as readily envisioned by one of ordinary skill in the art. Any of the foregoing nucleic acids or polypeptides can be tested by routine experimentation for retention of structural relation or activity to the nucleic acids and/or polypeptides disclosed herein.

The invention also provides isolated unique fragments of cancer associated antigen nucleic acid sequences or complements thereof. A unique fragment is one that is a 'signature' for the larger nucleic acid. It, for example, is long enough to assure that its precise sequence is not found in molecules within the human genome outside of the cancer associated antigen nucleic acids defined above (and human alleles). Those of ordinary skill in the art may apply no more than routine procedures to determine if a fragment is unique within the human genome. Unique fragments, however, exclude fragments completely composed of the nucleotide sequences of any of GenBank accession numbers listed in Table 4 or other previously published sequences as of the filing date of the priority documents for sequences listed in a respective priority document or the filing date of this application for sequences listed for the first time in this application which overlap the sequences of the invention.

A fragment which is completely composed of the sequence described in the foregoing

GenBank deposits is one which does not include any of the nucleotides unique to the sequences of the invention. Thus, a unique fragment must contain a nucleotide sequence other than the exact sequence of those in GenBank or fragments thereof. The difference may be an addition, deletion or substitution with respect to the GenBank sequence or it may be a sequence wholly separate from the GenBank sequence.

Unique fragments can be used as probes in Southern and Northern blot assays to identify such nucleic acids, or can be used in amplification assays such as those employing PCR. As known to those skilled in the art, large probes such as 200, 250, 300 or more nucleotides are preferred for certain uses such as Southern and Northern blots, while smaller fragments will be preferred for uses such as PCR. Unique fragments also can be used to produce fusion proteins for generating antibodies or determining binding of the polypeptide fragments, or for generating immunoassay components. Likewise, unique fragments can be employed to produce nonfused fragments of the cancer associated antigen polypeptides, useful, for example, in the preparation of antibodies, and in immunoassays. Unique fragments further can be used as antisense molecules to inhibit the expression of cancer associated antigen nucleic acids and polypeptides, particularly for therapeutic purposes as described in greater detail below.

As will be recognized by those skilled in the art, the size of the unique fragment will depend upon its conservancy in the genetic code. Thus, some regions of cancer associated antigen sequences and complements thereof will require longer segments to be unique while others will require only short segments, typically between 12 and 32 nucleotides (e.g. 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 and 32 or more bases long, up to the entire length of the disclosed sequence. As mentioned above, this disclosure intends to embrace each and every fragment of each sequence, beginning at the first nucleotide, the second nucleotide and so on, up to 8 nucleotides short of the end, and ending anywhere from nucleotide number 8, 9, 10 and so on for each sequence, up to the very last nucleotide (provided the sequence is unique as described above).

Virtually any segment of the polypeptide coding region of novel cancer associated antigen nucleic acids, or complements thereof, that is 18 or more nucleotides in length will be unique. Those skilled in the art are well versed in methods for selecting such sequences, typically on the basis of the ability of the unique fragment to selectively distinguish the sequence of interest from other sequences in the human genome of the fragment to those on known databases typically is all that is necessary, although *in vitro* confirmatory hybridization

and sequencing analysis may be performed.

Especially preferred include nucleic acids encoding a series of epitopes, known as "polytopes". The epitopes can be arranged in sequential or overlapping fashion (*see, e.g.,* Thomson et al., *Proc. Natl. Acad. Sci. USA* 92:5845-5849, 1995; Gilbert et al., *Nature Biotechnol.* 15:1280-1284, 1997), with or without the natural flanking sequences, and can be separated by unrelated linker sequences if desired. The polytope is processed to generate individual epitopes which are recognized by the immune system for generation of immune responses.

Thus, for example, peptides derived from a polypeptide having an amino acid sequence encoded by one of the nucleic acids disclosed herein, and which are presented by MHC molecules and recognized by CTL or T helper lymphocytes, can be combined with peptides from one or more other cancer associated antigens (e.g. by preparation of hybrid nucleic acids or polypeptides) to form "polytopes". The two or more peptides (or nucleic acids encoding the peptides) can be selected from those described herein, or they can include one or more peptides of previously known cancer associated antigens. Exemplary cancer associated peptide antigens that can be administered to induce or enhance an immune response are derived from tumor associated genes and encoded proteins including MAGE-A1, MAGE-A2, MAGE-A3, MAGE-A4, MAGE-A5, MAGE-A6, MAGE-A7, MAGE-A8, MAGE-A9, MAGE-A10, MAGE-A11, MAGE-A12, GAGE-1, GAGE-2, GAGE-3, GAGE-4, GAGE-5, GAGE-6, GAGE-7, GAGE-8, GAGE-9, BAGE-1, RAGE-1, LB33/MUM-1, PRAME, NAG, MAGE-B2, MAGE-B3, MAGE-B4, tyrosinase, brain glycogen phosphorylase, Melan-A, MAGE-C1, MAGE-C2, MAGE-C3, MAGE-C4, MAGE-C5, NY-ESO-1, LAGE-1, SSX-1, SSX-2 (HOM-MEL-40), SSX-4, SSX-5, SCP-1 and CT-7. See, for example, PCT application publication no. WO96/10577. Other examples will be known to one of ordinary skill in the art and can be used in the invention in a like manner as those disclosed herein. Other examples of HLA class I and HLA class II binding peptides will be known to one of ordinary skill in the art. For example, see the following references: Coulie, *Stem Cells* 13:393-403, 1995; Traversari et al., *J. Exp. Med.* 176:1453-1457, 1992; Chaux et al., *J. Immunol.* 163:2928-2936, 1999; Fujie et al., *Int. J. Cancer* 80:169-172, 1999; Tanzarella et al., *Cancer Res.* 59:2668-2674, 1999; van der Bruggen et al., *Eur. J. Immunol.* 24:2134-2140, 1994; Chaux et al., *J. Exp. Med.* 189:767-778, 1999; Kawashima et al., *Hum. Immunol.* 59:1-14, 1998; Tahara et al., *Clin. Cancer Res.* 5:2236-2241, 1999; Gaugler et al., *J. Exp. Med.* 179:921-930, 1994; van der Bruggen et al., *Eur. J. Immunol.* 24:3038-3043, 1994; Tanaka et

- al., *Cancer Res.* 57:4465-4468, 1997; Oiso et al., *Int. J. Cancer* 81:387-394, 1999; Herman et al., *Immunogenetics* 43:377-383, 1996; Manici et al., *J. Exp. Med.* 189:871-876, 1999; Duffour et al., *Eur. J. Immunol.* 29:3329-3337, 1999; Zorn et al., *Eur. J. Immunol.* 29:602-607, 1999; Huang et al., *J. Immunol.* 162:6849-6854, 1999; Boël et al., *Immunity* 2:167-175, 5 1995; Van den Eynde et al., *J. Exp. Med.* 182:689-698, 1995; De Backer et al., *Cancer Res.* 59:3157-3165, 1999; Jäger et al., *J. Exp. Med.* 187:265-270, 1998; Wang et al., *J. Immunol.* 161:3596-3606, 1998; Aarnoudse et al., *Int. J. Cancer* 82:442-448, 1999; Guilloux et al., *J. Exp. Med.* 183:1173-1183, 1996; Lupetti et al., *J. Exp. Med.* 188:1005-1016, 1998; Wölfel et al., *Eur. J. Immunol.* 24:759-764, 1994; Skipper et al., *J. Exp. Med.* 183:527-534, 1996; Kang 10 et al., *J. Immunol.* 155:1343-1348, 1995; Morel et al., *Int. J. Cancer* 83:755-759, 1999; Brichard et al., *Eur. J. Immunol.* 26:224-230, 1996; Kittlesen et al., *J. Immunol.* 160:2099-2106, 1998; Kawakami et al., *J. Immunol.* 161:6985-6992, 1998; Topalian et al., *J. Exp. Med.* 183:1965-1971, 1996; Kobayashi et al., *Cancer Research* 58:296-301, 1998; Kawakami et al., *J. Immunol.* 154:3961-3968, 1995; Tsai et al., *J. Immunol.* 158:1796-1802, 1997; Cox et al., 15 *Science* 264:716-719, 1994; Kawakami et al., *Proc. Natl. Acad. Sci. USA* 91:6458-6462, 1994; Skipper et al., *J. Immunol.* 157:5027-5033, 1996; Robbins et al., *J. Immunol.* 159:303-308, 1997; Castelli et al., *J. Immunol.* 162:1739-1748, 1999; Kawakami et al., *J. Exp. Med.* 180:347-352, 1994; Castelli et al., *J. Exp. Med.* 181:363-368, 1995; Schneider et al., *Int. J. Cancer* 75:451-458, 1998; Wang et al., *J. Exp. Med.* 183:1131-1140, 1996; Wang et al., *J. 20 Exp. Med.* 184:2207-2216, 1996; Parkhurst et al., *Cancer Research* 58:4895-4901, 1998; Tsang et al., *J. Natl Cancer Inst* 87:982-990, 1995; Correale et al., *J Natl Cancer Inst* 89:293-300, 1997; Coulie et al., *Proc. Natl. Acad. Sci. USA* 92:7976-7980, 1995; Wölfel et al., *Science* 269:1281-1284, 1995; Robbins et al., *J. Exp. Med.* 183:1185-1192, 1996; Brändle et al., *J. Exp. Med.* 183:2501-2508, 1996; ten Bosch et al., *Blood* 88:3522-3527, 1996; 25 Mandruzzato et al., *J. Exp. Med.* 186:785-793, 1997; Guéguen et al., *J. Immunol.* 160:6188-6194, 1998; Gjertsen et al., *Int. J. Cancer* 72:784-790, 1997; Gaudin et al., *J. Immunol.* 162:1730-1738, 1999; Chiari et al., *Cancer Res.* 59:5785-5792, 1999; Hogan et al., *Cancer Res.* 58:5144-5150, 1998; Pieper et al., *J. Exp. Med.* 189:757-765, 1999; Wang et al., *Science* 284:1351-1354, 1999; Fisk et al., *J. Exp. Med.* 181:2109-2117, 1995; Brossart et al., *Cancer 30 Res.* 58:732-736, 1998; Röpke et al., *Proc. Natl. Acad. Sci. USA* 93:14704-14707, 1996; Ikeda et al., *Immunity* 6:199-208, 1997; Ronsin et al., *J. Immunol.* 163:483-490, 1999; Vonderheide et al., *Immunity* 10:673-679, 1999.

One of ordinary skill in the art can prepare polypeptides comprising one or more

peptides and one or more of the foregoing cancer associated peptides, or nucleic acids encoding such polypeptides, according to standard procedures of molecular biology.

Thus polytopes are groups of two or more potentially immunogenic or immune response stimulating peptides which can be joined together in various arrangements (e.g. concatenated, overlapping). The polytope (or nucleic acid encoding the polytope) can be administered in a standard immunization protocol, e.g. to animals, to test the effectiveness of the polytope in stimulating, enhancing and/or provoking an immune response.

The peptides can be joined together directly or via the use of flanking sequences to form polytopes, and the use of polytopes as vaccines is well known in the art (see, e.g., Thomson et al., *Proc. Acad. Natl. Acad. Sci USA* 92(13):5845-5849, 1995; Gilbert et al., *Nature Biotechnol.* 15(12):1280-1284, 1997; Thomson et al., *J. Immunol.* 157(2):822-826, 1996; Tam et al., *J. Exp. Med.* 171(1):299-306, 1990). For example, Tam showed that polytopes consisting of both MHC class I and class II binding epitopes successfully generated antibody and protective immunity in a mouse model. Tam also demonstrated that polytopes comprising "strings" of epitopes are processed to yield individual epitopes which are presented by MHC molecules and recognized by CTLs. Thus polytopes containing various numbers and combinations of epitopes can be prepared and tested for recognition by CTLs and for efficacy in increasing an immune response.

It is known that tumors express a set of tumor antigens, of which only certain subsets may be expressed in the tumor of any given patient. Polytopes can be prepared which correspond to the different combination of epitopes representing the subset of tumor rejection antigens expressed in a particular patient. Polytopes also can be prepared to reflect a broader spectrum of tumor rejection antigens known to be expressed by a tumor type. Polytopes can be introduced to a patient in need of such treatment as polypeptide structures, or via the use of nucleic acid delivery systems known in the art (see, e.g., Allsopp et al., *Eur. J. Immunol.* 26(8):1951-1959, 1996). Adenovirus, pox virus, Ty-virus like particles, adeno-associated virus, plasmids, bacteria, etc. can be used in such delivery. One can test the polytope delivery systems in mouse models to determine efficacy of the delivery system. The systems also can be tested in human clinical trials.

In instances in which a human HLA class I molecule presents tumor rejection antigens derived from cancer associated nucleic acids, the expression vector may also include a nucleic acid sequence coding for the HLA molecule that presents any particular tumor rejection antigen derived from these nucleic acids and polypeptides. Alternatively, the nucleic acid

sequence coding for such a HLA molecule can be contained within a separate expression vector. In a situation where the vector contains both coding sequences, the single vector can be used to transfect a cell which does not normally express either one. Where the coding sequences for a cancer associated antigen precursor and the HLA molecule which presents it are contained on separate expression vectors, the expression vectors can be cotransfected. The cancer associated antigen precursor coding sequence may be used alone, when, e.g. the host cell already expresses a HLA molecule which presents a cancer associated antigen derived from precursor molecules. Of course, there is no limit on the particular host cell which can be used. As the vectors which contain the two coding sequences may be used in any antigen-presenting cells if desired, and the gene for cancer associated antigen precursor can be used in host cells which do not express a HLA molecule which presents a cancer associated antigen. Further, cell-free transcription systems may be used in lieu of cells.

As mentioned above, the invention embraces antisense oligonucleotides that selectively bind to a nucleic acid molecule encoding a cancer associated antigen polypeptide, to reduce the expression of cancer associated antigens. This is desirable in virtually any medical condition wherein a reduction of expression of cancer associated antigens is desirable, e.g., in the treatment of cancer. This is also useful for *in vitro* or *in vivo* testing of the effects of a reduction of expression of one or more cancer associated antigens.

As used herein, the term "antisense oligonucleotide" or "antisense" describes an oligonucleotide that is an oligoribonucleotide, oligodeoxyribonucleotide, modified oligoribonucleotide, or modified oligodeoxyribonucleotide which hybridizes under physiological conditions to DNA comprising a particular gene or to an mRNA transcript of that gene and, thereby, inhibits the transcription of that gene and/or the translation of that mRNA. The antisense molecules are designed so as to interfere with transcription or translation of a target gene upon hybridization with the target gene or transcript. Those skilled in the art will recognize that the exact length of the antisense oligonucleotide and its degree of complementarity with its target will depend upon the specific target selected, including the sequence of the target and the particular bases which comprise that sequence. It is preferred that the antisense oligonucleotide be constructed and arranged so as to bind selectively with the target under physiological conditions, i.e., to hybridize substantially more to the target sequence than to any other sequence in the target cell under physiological conditions. Based upon the sequences of nucleic acids encoding breast cancer associated antigen, or upon allelic or homologous genomic and/or cDNA sequences, one of skill in the

art can easily choose and synthesize any of a number of appropriate antisense molecules for use in accordance with the present invention. In order to be sufficiently selective and potent for inhibition, such antisense oligonucleotides should comprise at least 10 and, more preferably, at least 15 consecutive bases which are complementary to the target, although in certain cases modified oligonucleotides as short as 7 bases in length have been used successfully as antisense oligonucleotides (Wagner et al., *Nature Biotechnol.* 14:840-844, 1996). Most preferably, the antisense oligonucleotides comprise a complementary sequence of 20-30 bases. Although oligonucleotides may be chosen which are antisense to any region of the gene or mRNA transcripts, in preferred embodiments the antisense oligonucleotides correspond to N-terminal or 5' upstream sites such as translation initiation, transcription initiation or promoter sites. In addition, 3'-untranslated regions may be targeted. Targeting to mRNA splicing sites has also been used in the art but may be less preferred if alternative mRNA splicing occurs. In addition, the antisense is targeted, preferably, to sites in which mRNA secondary structure is not expected (see, e.g., Sainio et al., *Cell Mol. Neurobiol.* 14(5):439-457, 1994) and at which proteins are not expected to bind. Finally, although the listed sequences are cDNA sequences, one of ordinary skill in the art may easily derive the genomic DNA corresponding to the cDNA of a cancer associated antigen. Thus, the present invention also provides for antisense oligonucleotides which are complementary to the genomic DNA corresponding to nucleic acids encoding cancer associated antigens. Similarly, antisense to allelic or homologous cDNAs and genomic DNAs are enabled without undue experimentation.

In one set of embodiments, the antisense oligonucleotides of the invention may be composed of "natural" deoxyribonucleotides, ribonucleotides, or any combination thereof. That is, the 5' end of one native nucleotide and the 3' end of another native nucleotide may be covalently linked, as in natural systems, via a phosphodiester internucleoside linkage. These oligonucleotides may be prepared by art recognized methods which may be carried out manually or by an automated synthesizer. They also may be produced recombinantly by vectors.

In preferred embodiments, however, the antisense oligonucleotides of the invention also may include "modified" oligonucleotides. That is, the oligonucleotides may be modified in a number of ways which do not prevent them from hybridizing to their target but which enhance their stability or targeting or which otherwise enhance their therapeutic effectiveness.

The term "modified oligonucleotide" as used herein describes an oligonucleotide in

which (1) at least two of its nucleotides are covalently linked via a synthetic internucleoside linkage (i.e., a linkage other than a phosphodiester linkage between the 5' end of one nucleotide and the 3' end of another nucleotide) and/or (2) a chemical group not normally associated with nucleic acids has been covalently attached to the oligonucleotide. Preferred synthetic internucleoside linkages are phosphorothioates, alkylphosphonates, phosphorodithioates, phosphate esters, alkylphosphonothioates, phosphoramidates, carbamates, carbonates, phosphate triesters, acetamides, carboxymethyl esters and peptides.

The term "modified oligonucleotide" also encompasses oligonucleotides with a covalently modified base and/or sugar. For example, modified oligonucleotides include oligonucleotides having backbone sugars which are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 3' position and other than a phosphate group at the 5' position. Thus modified oligonucleotides may include a 2'-O-alkylated ribose group. In addition, modified oligonucleotides may include sugars such as arabinose instead of ribose. Base analogs such as C-5 propyne modified bases also can be included (*Nature Biotechnol.* 14:840-844, 1996). The present invention, thus, contemplates pharmaceutical preparations containing modified antisense molecules that are complementary to and hybridizable with, under physiological conditions, nucleic acids encoding the cancer associated antigen polypeptides, together with pharmaceutically acceptable carriers.

Antisense oligonucleotides may be administered as part of a pharmaceutical composition. Such a pharmaceutical composition may include the antisense oligonucleotides in combination with any standard physiologically and/or pharmaceutically acceptable carriers which are known in the art. The compositions should be sterile and contain a therapeutically effective amount of the antisense oligonucleotides in a unit of weight or volume suitable for administration to a patient. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. The term "physiologically acceptable" refers to a non-toxic material that is compatible with a biological system such as a cell, cell culture, tissue, or organism. The characteristics of the carrier will depend on the route of administration. Physiologically and pharmaceutically acceptable carriers include diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials which are well known in the art, as further described below.

As used herein, a "vector" may be any of a number of nucleic acids into which a desired sequence may be inserted by restriction and ligation for transport between different genetic environments or for expression in a host cell. Vectors are typically composed of DNA

although RNA vectors are also available. Vectors include, but are not limited to, plasmids, phagemids and virus genomes. A cloning vector is one which is able to replicate autonomously or integrated in the genome in a host cell, and which is further characterized by one or more endonuclease restriction sites at which the vector may be cut in a determinable fashion and into which a desired DNA sequence may be ligated such that the new recombinant vector retains its ability to replicate in the host cell. In the case of plasmids, replication of the desired sequence may occur many times as the plasmid increases in copy number within the host bacterium or just a single time per host before the host reproduces by mitosis. In the case of phage, replication may occur actively during a lytic phase or passively during a lysogenic phase. An expression vector is one into which a desired DNA sequence may be inserted by restriction and ligation such that it is operably joined to regulatory sequences and may be expressed as an RNA transcript. Vectors may further contain one or more marker sequences suitable for use in the identification of cells which have or have not been transformed or transfected with the vector. Markers include, for example, genes encoding proteins which increase or decrease either resistance or sensitivity to antibiotics or other compounds, genes which encode enzymes whose activities are detectable by standard assays known in the art (e.g.,  $\beta$ -galactosidase, luciferase or alkaline phosphatase), and genes which visibly affect the phenotype of transformed or transfected cells, hosts, colonies or plaques (e.g., green fluorescent protein). Preferred vectors are those capable of autonomous replication and expression of the structural gene products present in the DNA segments to which they are operably joined.

As used herein, a coding sequence and regulatory sequences are said to be “operably” joined when they are covalently linked in such a way as to place the expression or transcription of the coding sequence under the influence or control of the regulatory sequences. If it is desired that the coding sequences be translated into a functional protein, two DNA sequences are said to be operably joined if induction of a promoter in the 5’ regulatory sequences results in the transcription of the coding sequence and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region to direct the transcription of the coding sequences, or (3) interfere with the ability of the corresponding RNA transcript to be translated into a protein. Thus, a promoter region would be operably joined to a coding sequence if the promoter region were capable of effecting transcription of that DNA sequence such that the resulting transcript might be translated into the desired protein or polypeptide.

The precise nature of the regulatory sequences needed for gene expression may vary between species or cell types, but shall in general include, as necessary, 5' non-transcribed and 5' non-translated sequences involved with the initiation of transcription and translation respectively, such as a TATA box, capping sequence, CAAT sequence, and the like.

Especially, such 5' non-transcribed regulatory sequences will include a promoter region which includes a promoter sequence for transcriptional control of the operably joined gene. Regulatory sequences may also include enhancer sequences or upstream activator sequences as desired. The vectors of the invention may optionally include 5' leader or signal sequences. The choice and design of an appropriate vector is within the ability and discretion of one of ordinary skill in the art.

Expression vectors containing all the necessary elements for expression are commercially available and known to those skilled in the art. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, 1989. Cells are genetically engineered by the introduction into the cells of heterologous DNA (RNA) encoding a breast cancer associated antigen polypeptide or fragment or variant thereof. That heterologous DNA (RNA) is placed under operable control of transcriptional elements to permit the expression of the heterologous DNA in the host cell.

Preferred systems for mRNA expression in mammalian cells are those such as pRc/CMV (available from Invitrogen, Carlsbad, CA) that contain a selectable marker such as a gene that confers G418 resistance (which facilitates the selection of stably transfected cell lines) and the human cytomegalovirus (CMV) enhancer-promoter sequences. Additionally, suitable for expression in primate or canine cell lines is the pCEP4 vector (Invitrogen), which contains an Epstein Barr Virus (EBV) origin of replication, facilitating the maintenance of plasmid as a multicopy extrachromosomal element. Another expression vector is the pEF-BOS plasmid containing the promoter of polypeptide Elongation Factor 1 $\alpha$ , which stimulates efficiently transcription *in vitro*. The plasmid is described by Mishizuma and Nagata (*Nuc. Acids Res.* 18:5322, 1990), and its use in transfection experiments is disclosed by, for example, Demoulin (*Mol. Cell. Biol.* 16:4710-4716, 1996). Still another preferred expression vector is an adenovirus, described by Stratford-Perricaudet, which is defective for E1 and E3 proteins (*J. Clin. Invest.* 90:626-630, 1992). The use of the adenovirus as an Adeno.P1A recombinant for the expression of an antigen is disclosed by Warnier et al., in intradermal injection in mice for immunization against P1A (*Int. J. Cancer*, 67:303-310, 1996). Additional vectors for delivery of nucleic acid are provided below.

The invention also embraces so-called expression kits, which allow the artisan to prepare a desired expression vector or vectors. Such expression kits include at least separate portions of a vector and one or more of the previously discussed cancer associated antigen nucleic acid molecules. Other components may be added, as desired, as long as the previously mentioned nucleic acid molecules, which are required, are included. The invention also includes kits for amplification of a cancer associated antigen nucleic acid, including at least one pair of amplification primers which hybridize to a cancer associated antigen nucleic acid. The primers preferably are 12-32 nucleotides in length and are non-overlapping to prevent formation of "primer-dimers". One of the primers will hybridize to one strand of the cancer associated antigen nucleic acid and the second primer will hybridize to the complementary strand of the cancer associated antigen nucleic acid, in an arrangement which permits amplification of the cancer associated antigen nucleic acid. Selection of appropriate primer pairs is standard in the art. For example, the selection can be made with assistance of a computer program designed for such a purpose, optionally followed by testing the primers for amplification specificity and efficiency.

The invention also permits the construction of cancer associated antigen gene "knock-outs" in cells and in animals, providing materials for studying certain aspects of cancer and immune system responses to cancer.

The invention also provides isolated polypeptides (including whole proteins and partial proteins) encoded by the foregoing cancer associated antigen nucleic acids. Such polypeptides are useful, for example, alone or as fusion proteins to generate antibodies, as components of an immunoassay or diagnostic assay or as therapeutics. Cancer associated antigen polypeptides can be isolated from biological samples including tissue or cell homogenates, and can also be expressed recombinantly in a variety of prokaryotic and eukaryotic expression systems by constructing an expression vector appropriate to the expression system, introducing the expression vector into the expression system, and isolating the recombinantly expressed protein. Short polypeptides, including antigenic peptides (such as are presented by MHC molecules on the surface of a cell for immune recognition) also can be synthesized chemically using well-established methods of peptide synthesis.

A unique fragment of a cancer associated antigen polypeptide, in general, has the features and characteristics of unique fragments as discussed above in connection with nucleic acids. As will be recognized by those skilled in the art, the size of the unique fragment will depend upon factors such as whether the fragment constitutes a portion of a conserved protein

domain. Thus, some regions of cancer associated antigens will require longer segments to be unique while others will require only short segments, typically between 5 and 12 amino acids (e.g. 5, 6, 7, 8, 9, 10, 11 or 12 or more amino acids including each integer up to the full length).

5 Unique fragments of a polypeptide preferably are those fragments which retain a distinct functional capability of the polypeptide. Functional capabilities which can be retained in a unique fragment of a polypeptide include interaction with antibodies, interaction with other polypeptides or fragments thereof, selective binding of nucleic acids or proteins, and enzymatic activity. One important activity is the ability to act as a signature for identifying  
10 the polypeptide. Another is the ability to complex with HLA and to provoke in a human an immune response. Those skilled in the art are well versed in methods for selecting unique amino acid sequences, typically on the basis of the ability of the unique fragment to selectively distinguish the sequence of interest from non-family members. A comparison of the sequence of the fragment to those on known databases typically is all that is necessary.

15 The invention embraces variants of the cancer associated antigen polypeptides described above. As used herein, a "variant" of a cancer associated antigen polypeptide is a polypeptide which contains one or more modifications to the primary amino acid sequence of a cancer associated antigen polypeptide. Modifications which create a cancer associated antigen variant can be made to a cancer associated antigen polypeptide 1) to reduce or  
20 eliminate an activity of a cancer associated antigen polypeptide; 2) to enhance a property of a cancer associated antigen polypeptide, such as protein stability in an expression system or the stability of protein-protein binding; 3) to provide a novel activity or property to a cancer associated antigen polypeptide, such as addition of an antigenic epitope or addition of a detectable moiety; or 4) to provide equivalent or better binding to an HLA molecule.

25 Modifications to a cancer associated antigen polypeptide are typically made to the nucleic acid which encodes the cancer associated antigen polypeptide, and can include deletions, point mutations, truncations, amino acid substitutions and additions of amino acids or non-amino acid moieties. Alternatively, modifications can be made directly to the polypeptide, such as by cleavage, addition of a linker molecule, addition of a detectable moiety, such as  
30 biotin, addition of a fatty acid, and the like. Modifications also embrace fusion proteins comprising all or part of the cancer associated antigen amino acid sequence. One of skill in the art will be familiar with methods for predicting the effect on protein conformation of a change in protein sequence, and can thus "design" a variant cancer associated antigen

polypeptide according to known methods. One example of such a method is described by Dahiyat and Mayo in *Science* 278:82-87, 1997, whereby proteins can be designed *de novo*. The method can be applied to a known protein to vary a only a portion of the polypeptide sequence. By applying the computational methods of Dahiyat and Mayo, specific variants of  
5 a cancer associated antigen polypeptide can be proposed and tested to determine whether the variant retains a desired conformation.

In general, variants include cancer associated antigen polypeptides which are modified specifically to alter a feature of the polypeptide unrelated to its desired physiological activity.

For example, cysteine residues can be substituted or deleted to prevent unwanted disulfide  
10 linkages. Similarly, certain amino acids can be changed to enhance expression of a breast cancer associated antigen polypeptide by eliminating proteolysis by proteases in an expression system (e.g., dibasic amino acid residues in yeast expression systems in which KEX2 protease activity is present).

Mutations of a nucleic acid which encode a cancer associated antigen polypeptide  
15 preferably preserve the amino acid reading frame of the coding sequence, and preferably do not create regions in the nucleic acid which are likely to hybridize to form secondary structures, such a hairpins or loops, which can be deleterious to expression of the variant polypeptide.

Mutations can be made by selecting an amino acid substitution, or by random  
20 mutagenesis of a selected site in a nucleic acid which encodes the polypeptide. Variant polypeptides are then expressed and tested for one or more activities to determine which mutation provides a variant polypeptide with the desired properties. Further mutations can be made to variants (or to non-variant cancer associated antigen polypeptides) which are silent as to the amino acid sequence of the polypeptide, but which provide preferred codons for  
25 translation in a particular host. The preferred codons for translation of a nucleic acid in, e.g., *E. coli*, are well known to those of ordinary skill in the art. Still other mutations can be made to the noncoding sequences of a cancer associated antigen gene or cDNA clone to enhance expression of the polypeptide. The activity of variants of cancer associated antigen polypeptides can be tested by cloning the gene encoding the variant cancer associated antigen  
30 polypeptide into a bacterial or mammalian expression vector, introducing the vector into an appropriate host cell, expressing the variant cancer associated antigen polypeptide, and testing for a functional capability of the cancer associated antigen polypeptides as disclosed herein. For example, the variant cancer associated antigen polypeptide can be tested for reaction with

autologous or allogeneic sera as disclosed in the Examples. Preparation of other variant polypeptides may favor testing of other activities, as will be known to one of ordinary skill in the art.

The skilled artisan will also realize that conservative amino acid substitutions may be made in cancer associated antigen polypeptides to provide functionally equivalent variants of the foregoing polypeptides, i.e, the variants retain the functional capabilities of the cancer associated antigen polypeptides. As used herein, a "conservative amino acid substitution" refers to an amino acid substitution which does not alter the relative charge or size characteristics of the protein in which the amino acid substitution is made. Variants can be prepared according to methods for altering polypeptide sequence known to one of ordinary skill in the art such as are found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. Exemplary functionally equivalent variants of the cancer associated antigen polypeptides include conservative amino acid substitutions of in the amino acid sequences of proteins disclosed herein. Conservative substitutions of amino acids include substitutions made amongst amino acids within the following groups: (a) M, I, L, V; (b) F, Y, W; (c) K, R, H; (d) A, G; (e) S, T; (f) Q, N; and (g) E, D.

For example, upon determining that a peptide derived from a cancer associated antigen polypeptide is presented by an MHC molecule and recognized by CTLs (e.g., as described in the Examples), one can make conservative amino acid substitutions to the amino acid sequence of the peptide, particularly at residues which are thought not to be direct contact points with the MHC molecule. For example, methods for identifying functional variants of HLA class II binding peptides are provided in a published PCT application of Strominger and Wucherpfennig (PCT/US96/03182). Peptides bearing one or more amino acid substitutions also can be tested for concordance with known HLA/MHC motifs prior to synthesis using, e.g. the computer program described by D'Amaro and Drijfhout (D'Amaro et al., *Human Immunol.* 43:13-18, 1995; Drijfhout et al., *Human Immunol.* 43:1-12, 1995). The substituted peptides can then be tested for binding to the MHC molecule and recognition by CTLs when bound to MHC. These variants can be tested for improved stability and are useful, *inter alia*, in vaccine compositions.

Conservative amino-acid substitutions in the amino acid sequence of cancer associated

antigen polypeptides to produce functionally equivalent variants of cancer associated antigen polypeptides typically are made by alteration of a nucleic acid encoding a cancer associated antigen polypeptide. Such substitutions can be made by a variety of methods known to one of ordinary skill in the art. For example, amino acid substitutions may be made by PCR-directed mutation, site-directed mutagenesis according to the method of Kunkel (Kunkel, *Proc. Nat. Acad. Sci. U.S.A.* 82: 488-492, 1985), or by chemical synthesis of a gene encoding a cancer associated antigen polypeptide. Where amino acid substitutions are made to a small unique fragment of a cancer associated antigen polypeptide, such as an antigenic epitope recognized by autologous or allogeneic sera or cytolytic T lymphocytes, the substitutions can be made by directly synthesizing the peptide. The activity of functionally equivalent fragments of cancer associated antigen polypeptides can be tested by cloning the gene encoding the altered cancer associated antigen polypeptide into a bacterial or mammalian expression vector, introducing the vector into an appropriate host cell, expressing the altered cancer associated antigen polypeptide, and testing for a functional capability of the cancer associated antigen polypeptides as disclosed herein. Peptides which are chemically synthesized can be tested directly for function, e.g., for binding to antisera recognizing associated antigens.

The invention as described herein has a number of uses, some of which are described elsewhere herein. First, the invention permits isolation of the cancer associated antigen protein molecules. A variety of methodologies well-known to the skilled practitioner can be utilized to obtain isolated cancer associated antigen molecules. The polypeptide may be purified from cells which naturally produce the polypeptide by chromatographic means or immunological recognition. Alternatively, an expression vector may be introduced into cells to cause production of the polypeptide. In another method, mRNA transcripts may be microinjected or otherwise introduced into cells to cause production of the encoded polypeptide. Translation of mRNA in cell-free extracts such as the reticulocyte lysate system also may be used to produce polypeptide. Those skilled in the art also can readily follow known methods for isolating cancer associated antigen polypeptides. These include, but are not limited to, immunochromatography, HPLC, size-exclusion chromatography, ion-exchange chromatography and immune-affinity chromatography.

The isolation and identification of cancer associated antigen genes also makes it possible for the artisan to diagnose a disorder characterized by expression of cancer associated antigens. These methods involve determining expression of one or more cancer associated antigen nucleic acids, and/or encoded cancer associated antigen polypeptides and/or peptides

derived therefrom. In the former situation, such determinations can be carried out via any standard nucleic acid determination assay, including the polymerase chain reaction, or assaying with labeled hybridization probes. In the latter situation, such determinations can be carried out by screening patient antisera for recognition of the polypeptide.

5           The invention also makes it possible isolate proteins which bind to cancer associated antigens as disclosed herein, including antibodies and cellular binding partners of the cancer associated antigens. Additional uses are described further herein.

          The invention also provides, in certain embodiments, "dominant negative" polypeptides derived from cancer associated antigen polypeptides. A dominant negative  
10   polypeptide is an inactive variant of a protein, which, by interacting with the cellular machinery, displaces an active protein from its interaction with the cellular machinery or competes with the active protein, thereby reducing the effect of the active protein. For example, a dominant negative receptor which binds a ligand but does not transmit a signal in response to binding of the ligand can reduce the biological effect of expression of the ligand.  
15   Likewise, a dominant negative catalytically-inactive kinase which interacts normally with target proteins but does not phosphorylate the target proteins can reduce phosphorylation of the target proteins in response to a cellular signal. Similarly, a dominant negative transcription factor which binds to a promoter site in the control region of a gene but does not increase gene transcription can reduce the effect of a normal transcription factor by occupying  
20   promoter binding sites without increasing transcription.

          The end result of the expression of a dominant negative polypeptide in a cell is a reduction in function of active proteins. One of ordinary skill in the art can assess the potential for a dominant negative variant of a protein, and using standard mutagenesis techniques to create one or more dominant negative variant polypeptides. For example, given  
25   the teachings contained herein of small cell lung cancer associated antigens, especially those which are similar to known proteins which have known activities, one of ordinary skill in the art can modify the sequence of the cancer associated antigens by site-specific mutagenesis, scanning mutagenesis, partial gene deletion or truncation, and the like. See, e.g., U.S. Patent No. 5,580,723 and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second  
30   Edition, Cold Spring Harbor Laboratory Press, 1989. The skilled artisan then can test the population of mutagenized polypeptides for diminution in a selected and/or for retention of such an activity. Other similar methods for creating and testing dominant negative variants of a protein will be apparent to one of ordinary skill in the art.

The invention also involves agents such as polypeptides which bind to cancer associated antigen polypeptides. Such binding agents can be used, for example, in screening assays to detect the presence or absence of cancer associated antigen polypeptides and complexes of cancer associated antigen polypeptides and their binding partners and in purification protocols to isolated cancer associated antigen polypeptides and complexes of cancer associated antigen polypeptides and their binding partners. Such agents also can be used to inhibit the native activity of the cancer associated antigen polypeptides, for example, by binding to such polypeptides.

The invention, therefore, embraces peptide binding agents which, for example, can be antibodies or fragments of antibodies having the ability to selectively bind to cancer associated antigen polypeptides. Antibodies include polyclonal and monoclonal antibodies, prepared according to conventional methodology.

Significantly, as is well-known in the art, only a small portion of an antibody molecule, the paratope, is involved in the binding of the antibody to its epitope (see, in general, Clark, W.R. (1986) The Experimental Foundations of Modern Immunology Wiley & Sons, Inc., New York; Roitt, I. (1991) Essential Immunology, 7th Ed., Blackwell Scientific Publications, Oxford). The pFc' and Fc regions, for example, are effectors of the complement cascade but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically cleaved, or which has been produced without the pFc' region, designated an F(ab')<sub>2</sub> fragment, retains both of the antigen binding sites of an intact antibody. Similarly, an antibody from which the Fc region has been enzymatically cleaved, or which has been produced without the Fc region, designated an Fab fragment, retains one of the antigen binding sites of an intact antibody molecule. Proceeding further, Fab fragments consist of a covalently bound antibody light chain and a portion of the antibody heavy chain denoted Fd. The Fd fragments are the major determinant of antibody specificity (a single Fd fragment may be associated with up to ten different light chains without altering antibody specificity) and Fd fragments retain epitope-binding ability in isolation.

Within the antigen-binding portion of an antibody, as is well-known in the art, there are complementarity determining regions (CDRs), which directly interact with the epitope of the antigen, and framework regions (FRs), which maintain the tertiary structure of the paratope (see, in general, Clark, 1986; Roitt, 1991). In both the heavy chain Fd fragment and the light chain of IgG immunoglobulins, there are four framework regions (FR1 through FR4) separated respectively by three complementarity determining regions (CDR1 through CDR3).

The CDRs, and in particular the CDR3 regions, and more particularly the heavy chain CDR3, are largely responsible for antibody specificity.

It is now well-established in the art that the non-CDR regions of a mammalian antibody may be replaced with similar regions of conspecific or heterospecific antibodies while retaining the epitopic specificity of the original antibody. This is most clearly manifested in the development and use of "humanized" antibodies in which non-human CDRs are covalently joined to human FR and/or Fc/pFc' regions to produce a functional antibody. See, e.g., U.S. patents 4,816,567, 5,225,539, 5,585,089, 5,693,762 and 5,859,205.

Thus, for example, PCT International Publication Number WO 92/04381 teaches the production and use of humanized murine RSV antibodies in which at least a portion of the murine FR regions have been replaced by FR regions of human origin. Such antibodies, including fragments of intact antibodies with antigen-binding ability, are often referred to as "chimeric" antibodies.

Thus, as will be apparent to one of ordinary skill in the art, the present invention also provides for F(ab')<sub>2</sub>, Fab, Fv and Fd fragments; chimeric antibodies in which the Fc and/or FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric F(ab')<sub>2</sub> fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric Fab fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; and chimeric Fd fragment antibodies in which the FR and/or CDR1 and/or CDR2 regions have been replaced by homologous human or non-human sequences. The present invention also includes so-called single chain antibodies.

Thus, the invention involves polypeptides of numerous size and type that bind specifically to cancer associated antigen polypeptides, and complexes of both cancer associated antigen polypeptides and their binding partners. These polypeptides may be derived also from sources other than antibody technology. For example, such polypeptide binding agents can be provided by degenerate peptide libraries which can be readily prepared in solution, in immobilized form or as phage display libraries. Combinatorial libraries also can be synthesized of peptides containing one or more amino acids. Libraries further can be synthesized of peptoids and non-peptide synthetic moieties.

Phage display can be particularly effective in identifying binding peptides useful according to the invention. Briefly, one prepares a phage library (using e.g. m13, fd, or

lambda phage), displaying inserts from 4 to about 80 amino acid residues using conventional procedures. The inserts may represent, for example, a completely degenerate or biased array. One then can select phage-bearing inserts which bind to the cancer associated antigen polypeptide. This process can be repeated through several cycles of reselection of phage that bind to the cancer associated antigen polypeptide. Repeated rounds lead to enrichment of phage bearing particular sequences. DNA sequence analysis can be conducted to identify the sequences of the expressed polypeptides. The minimal linear portion of the sequence that binds to the cancer associated antigen polypeptide can be determined. One can repeat the procedure using a biased library containing inserts containing part or all of the minimal linear portion plus one or more additional degenerate residues upstream or downstream thereof. Yeast two-hybrid screening methods also may be used to identify polypeptides that bind to the cancer associated antigen polypeptides. Thus, the cancer associated antigen polypeptides of the invention, or a fragment thereof, can be used to screen peptide libraries, including phage display libraries, to identify and select peptide binding partners of the cancer associated antigen polypeptides of the invention. Such molecules can be used, as described, for screening assays, for purification protocols, for interfering directly with the functioning of cancer associated antigen and for other purposes that will be apparent to those of ordinary skill in the art.

As detailed herein, the foregoing antibodies and other binding molecules may be used for example to identify tissues expressing protein or to purify protein. Antibodies also may be coupled to specific diagnostic labeling agents for imaging of cells and tissues that express cancer associated antigens or to therapeutically useful agents according to standard coupling procedures. Diagnostic agents include, but are not limited to, barium sulfate, iocetamic acid, iopanoic acid, ipodate calcium, diatrizoate sodium, diatrizoate meglumine, metrizamide, tyropanoate sodium and radiodiagnostics including positron emitters such as fluorine-18 and carbon-11, gamma emitters such as iodine-123, technitium-99m, iodine-131 and indium-111, nuclides for nuclear magnetic resonance such as fluorine and gadolinium. Other diagnostic agents useful in the invention will be apparent to one of ordinary skill in the art. As used herein, "therapeutically useful agents" include any therapeutic molecule which desirably is targeted selectively to a cell expressing one of the cancer antigens disclosed herein, including antineoplastic agents, radioiodinated compounds, toxins, other cytostatic or cytolytic drugs, and so forth. Antineoplastic therapeutics are well known and include: aminoglutethimide, azathioprine, bleomycin sulfate, busulfan, carmustine, chlorambucil, cisplatin,

cyclophosphamide, cyclosporine, cytarabidine, dacarbazine, dactinomycin, daunorubicin, doxorubicin, taxol, etoposide, fluorouracil, interferon- $\alpha$ , lomustine, mercaptopurine, methotrexate, mitotane, procarbazine HCl, thioguanine, vinblastine sulfate and vincristine sulfate. Additional antineoplastic agents include those disclosed in Chapter 52,

Antineoplastic Agents (Paul Calabresi and Bruce A. Chabner), and the introduction thereto, 1202-1263, of Goodman and Gilman's "The Pharmacological Basis of Therapeutics", Eighth Edition, 1990, McGraw-Hill, Inc. (Health Professions Division). Toxins can be proteins such as, for example, pokeweed anti-viral protein, cholera toxin, pertussis toxin, ricin, gelonin, abrin, diphtheria exotoxin, or *Pseudomonas* exotoxin. Toxin moieties can also be high energy-emitting radionuclides such as cobalt-60.

In the foregoing methods, antibodies prepared according to the invention also preferably are specific for the small cell lung cancer associated antigen/MHC complexes described herein.

When "disorder" is used herein, it refers to any pathological condition where the cancer associated antigens are expressed. An example of such a disorder is cancer, with lung cancers including small cell lung cancer and non-small cell lung cancer, melanoma, colon cancer, breast cancer, head and neck cancer, transitional cancer, leiomyosarcoma and synovial sarcoma as particular examples.

Samples of tissue and/or cells for use in the various methods described herein can be obtained through standard methods such as tissue biopsy, including punch biopsy and cell scraping, and collection of blood or other bodily fluids by aspiration or other methods.

In certain embodiments of the invention, an immunoreactive cell sample is removed from a subject. By "immunoreactive cell" is meant a cell which can mature into an immune cell (such as a B cell, a helper T cell, or a cytolytic T cell) upon appropriate stimulation. Thus immunoreactive cells include CD34<sup>+</sup> hematopoietic stem cells, immature T cells and immature B cells. When it is desired to produce cytolytic T cells which recognize a cancer associated antigen, the immunoreactive cell is contacted with a cell which expresses a cancer associated antigen under conditions favoring production, differentiation and/or selection of cytolytic T cells; the differentiation of the T cell precursor into a cytolytic T cell upon exposure to antigen is similar to clonal selection of the immune system.

Some therapeutic approaches based upon the disclosure are premised on a response by a subject's immune system, leading to lysis of antigen presenting cells, such as breast cancer cells which present one or more cancer associated antigens. One such approach is the

administration of autologous CTLs specific to a cancer associated antigen/MHC complex to a subject with abnormal cells of the phenotype at issue. It is within the ability of one of ordinary skill in the art to develop such CTLs *in vitro*. An example of a method for T cell differentiation is presented in International Application number PCT/US96/05607. Generally, a sample of cells taken from a subject, such as blood cells, are contacted with a cell presenting the complex and capable of provoking CTLs to proliferate. The target cell can be a transfectant, such as a COS cell. These transfectants present the desired complex of their surface and, when combined with a CTL of interest, stimulate its proliferation. COS cells are widely available, as are other suitable host cells. Specific production of CTL clones is well known in the art. The clonally expanded autologous CTLs then are administered to the subject.

Another method for selecting antigen-specific CTL clones has recently been described (Altman et al., *Science* 274:94-96, 1996; Dunbar et al., *Curr. Biol.* 8:413-416, 1998), in which fluorogenic tetramers of MHC class I molecule/peptide complexes are used to detect specific CTL clones. Briefly, soluble MHC class I molecules are folded *in vitro* in the presence of  $\beta_2$ -microglobulin and a peptide antigen which binds the class I molecule. After purification, the MHC/peptide complex is purified and labeled with biotin. Tetramers are formed by mixing the biotinylated peptide-MHC complex with labeled avidin (e.g. phycoerythrin) at a molar ratio of 4:1. Tetramers are then contacted with a source of CTLs such as peripheral blood or lymph node. The tetramers bind CTLs which recognize the peptide antigen/MHC class I complex. Cells bound by the tetramers can be sorted by fluorescence activated cell sorting to isolate the reactive CTLs. The isolated CTLs then can be expanded *in vitro* for use as described herein.

To detail a therapeutic methodology, referred to as adoptive transfer (Greenberg, *J. Immunol.* 136(5): 1917, 1986; Riddel et al., *Science* 257: 238, 1992; Lynch et al, *Eur. J. Immunol.* 21: 1403-1410, 1991; Kast et al., *Cell* 59: 603-614, 1989), cells presenting the desired complex (e.g., dendritic cells) are combined with CTLs leading to proliferation of the CTLs specific thereto. The proliferated CTLs are then administered to a subject with a cellular abnormality which is characterized by certain of the abnormal cells presenting the particular complex. The CTLs then lyse the abnormal cells, thereby achieving the desired therapeutic goal.

The foregoing therapy assumes that at least some of the subject's abnormal cells present the relevant HLA/cancer associated antigen complex. This can be determined very

easily, as the art is very familiar with methods for identifying cells which present a particular HLA molecule, as well as how to identify cells expressing DNA of the pertinent sequences, in this case a cancer associated antigen sequence. Once cells presenting the relevant complex are identified via the foregoing screening methodology, they can be combined with a sample from a patient, where the sample contains CTLs. If the complex presenting cells are lysed by the mixed CTL sample, then it can be assumed that a cancer associated antigen is being presented, and the subject is an appropriate candidate for the therapeutic approaches set forth *supra*.

Adoptive transfer is not the only form of therapy that is available in accordance with the invention. CTLs can also be provoked *in vivo*, using a number of approaches. One approach is the use of non-proliferative cells expressing the complex. The cells used in this approach may be those that normally express the complex, such as irradiated tumor cells or cells transfected with one or both of the genes necessary for presentation of the complex (i.e. the antigenic peptide and the presenting HLA molecule). Chen et al. (*Proc. Natl. Acad. Sci. USA* 88: 110-114,1991) exemplifies this approach, showing the use of transfected cells expressing HPVE7 peptides in a therapeutic regime. Various cell types may be used. Similarly, vectors carrying one or both of the genes of interest may be used. Viral or bacterial vectors are especially preferred. For example, nucleic acids which encode a cancer associated antigen polypeptide or peptide may be operably linked to promoter and enhancer sequences which direct expression of the cancer associated antigen polypeptide or peptide in certain tissues or cell types. The nucleic acid may be incorporated into an expression vector. Expression vectors may be unmodified extrachromosomal nucleic acids, plasmids or viral genomes constructed or modified to enable insertion of exogenous nucleic acids, such as those encoding cancer associated antigen, as described elsewhere herein. Nucleic acids encoding a cancer associated antigen also may be inserted into a retroviral genome, thereby facilitating integration of the nucleic acid into the genome of the target tissue or cell type. In these systems, the gene of interest is carried by a microorganism, e.g., a Vaccinia virus, pox virus, herpes simplex virus, retrovirus or adenovirus, and the materials de facto "infect" host cells. The cells which result present the complex of interest, and are recognized by autologous CTLs, which then proliferate.

A similar effect can be achieved by combining the cancer associated antigen or a stimulatory fragment thereof with an adjuvant to facilitate incorporation into antigen presenting cells *in vivo*. The cancer associated antigen polypeptide is processed to yield the peptide partner of the HLA molecule while a cancer associated antigen peptide may be

presented without the need for further processing. Generally, subjects can receive an intradermal injection of an effective amount of the cancer associated antigen. Initial doses can be followed by booster doses, following immunization protocols standard in the art. Preferred cancer associated antigens include those found to react with allogeneic cancer antisera, shown in the examples below.

The invention involves the use of various materials disclosed herein to “immunize” subjects or as “vaccines”. As used herein, “immunization” or “vaccination” means increasing or activating an immune response against an antigen. It does not require elimination or eradication of a condition but rather contemplates the clinically favorable enhancement of an immune response toward an antigen. Generally accepted animal models can be used for testing of immunization against cancer using a cancer associated antigen nucleic acid. For example, human cancer cells can be introduced into a mouse to create a tumor, and one or more cancer associated antigen nucleic acids can be delivered by the methods described herein. The effect on the cancer cells (e.g., reduction of tumor size) can be assessed as a measure of the effectiveness of the cancer associated antigen nucleic acid immunization. Of course, testing of the foregoing animal model using more conventional methods for immunization include the administration of one or more cancer associated antigen polypeptides or peptides derived therefrom, optionally combined with one or more adjuvants and/or cytokines to boost the immune response. Methods for immunization, including formulation of a vaccine composition and selection of doses, route of administration and the schedule of administration (e.g. primary and one or more booster doses), are well known in the art. The tests also can be performed in humans, where the end point is to test for the presence of enhanced levels of circulating CTLs against cells bearing the antigen, to test for levels of circulating antibodies against the antigen, to test for the presence of cells expressing the antigen and so forth.

As part of the immunization compositions, one or more cancer associated antigens or stimulatory fragments thereof are administered with one or more adjuvants to induce an immune response or to increase an immune response. An adjuvant is a substance incorporated into or administered with antigen which potentiates the immune response. Adjuvants may enhance the immunological response by providing a reservoir of antigen (extracellularly or within macrophages), activating macrophages and stimulating specific sets of lymphocytes. Adjuvants of many kinds are well known in the art. Specific examples of adjuvants include monophosphoryl lipid A (MPL, SmithKline Beecham), a congener obtained after purification

and acid hydrolysis of *Salmonella minnesota* Re 595 lipopolysaccharide; saponins including QS21 (SmithKline Beecham), a pure QA-21 saponin purified from *Quillja saponaria* extract; DQS21, described in PCT application WO96/33739 (SmithKline Beecham); QS-7, QS-17, QS-18, and QS-L1 (So et al., *Mol. Cells* 7:178-186, 1997); incomplete Freund's adjuvant; complete Freund's adjuvant; montanide; and various water-in-oil emulsions prepared from biodegradable oils such as squalene and/or tocopherol. Preferably, the peptides are administered mixed with a combination of DQS21/MPL. The ratio of DQS21 to MPL typically will be about 1:10 to 10:1, preferably about 1:5 to 5:1 and more preferably about 1:1. Typically for human administration, DQS21 and MPL will be present in a vaccine formulation in the range of about 1 µg to about 100 µg. Other adjuvants are known in the art and can be used in the invention (*see, e.g. Goding, Monoclonal Antibodies: Principles and Practice*, 2nd Ed., 1986). Methods for the preparation of mixtures or emulsions of peptide and adjuvant are well known to those of skill in the art of vaccination.

Other agents which stimulate the immune response of the subject can also be administered to the subject. For example, other cytokines are also useful in vaccination protocols as a result of their lymphocyte regulatory properties. Many other cytokines useful for such purposes will be known to one of ordinary skill in the art, including interleukin-12 (IL-12) which has been shown to enhance the protective effects of vaccines (*see, e.g., Science* 268: 1432-1434, 1995), GM-CSF and IL-18. Thus cytokines can be administered in conjunction with antigens and adjuvants to increase the immune response to the antigens.

There are a number of immune response potentiating compounds that can be used in vaccination protocols. These include costimulatory molecules provided in either protein or nucleic acid form. Such costimulatory molecules include the B7-1 and B7-2 (CD80 and CD86 respectively) molecules which are expressed on dendritic cells (DC) and interact with the CD28 molecule expressed on the T cell. This interaction provides costimulation (signal 2) to an antigen/MHC/TCR stimulated (signal 1) T cell, increasing T cell proliferation and effector function. B7 also interacts with CTLA4 (CD152) on T cells and studies involving CTLA4 and B7 ligands indicate that the B7-CTLA4 interaction can enhance antitumor immunity and CTL proliferation (Zheng P., et al. *Proc. Natl. Acad. Sci. USA* 95 (11):6284-6289 (1998)).

B7 typically is not expressed on tumor cells so they are not efficient antigen presenting cells (APCs) for T cells. Induction of B7 expression would enable the tumor cells to stimulate more efficiently CTL proliferation and effector function. A combination of B7/IL-6/IL-12 costimulation has been shown to induce IFN-gamma and a Th1 cytokine profile in the T cell

population leading to further enhanced T cell activity (Gajewski et al., *J. Immunol.*, 154:5637-5648 (1995)). Tumor cell transfection with B7 has been discussed in relation to *in vitro* CTL expansion for adoptive transfer immunotherapy by Wang et al., (*J. Immunol.*, 19:1-8 (1986)). Other delivery mechanisms for the B7 molecule would include nucleic acid (naked DNA) immunization (Kim J., et al. *Nat Biotechnol.*, 15:7:641-646 (1997)) and recombinant viruses such as adeno and pox (Wendtnier et al., *Gene Ther.*, 4:7:726-735 (1997)). These systems are all amenable to the construction and use of expression cassettes for the coexpression of B7 with other molecules of choice such as the antigens or fragment(s) of antigens discussed herein (including polytopes) or cytokines. These delivery systems can be used for induction of the appropriate molecules *in vitro* and for *in vivo* vaccination situations. The use of anti-CD28 antibodies to directly stimulate T cells *in vitro* and *in vivo* could also be considered. Similarly, the inducible co-stimulatory molecule ICOS which induces T cell responses to foreign antigen could be modulated, for example, by use of anti-ICOS antibodies (Hutloff et al., *Nature* 397:263-266, 1999).

Lymphocyte function associated antigen-3 (LFA-3) is expressed on APCs and some tumor cells and interacts with CD2 expressed on T cells. This interaction induces T cell IL-2 and IFN-gamma production and can thus complement but not substitute, the B7/CD28 costimulatory interaction (Parra et al., *J. Immunol.*, 158:637-642 (1997), Fenton et al., *J. Immunother.*, 21:2:95-108 (1998)).

Lymphocyte function associated antigen-1 (LFA-1) is expressed on leukocytes and interacts with ICAM-1 expressed on APCs and some tumor cells. This interaction induces T cell IL-2 and IFN-gamma production and can thus complement but not substitute, the B7/CD28 costimulatory interaction (Fenton et al., *J. Immunother.*, 21:2:95-108 (1998)). LFA-1 is thus a further example of a costimulatory molecule that could be provided in a vaccination protocol in the various ways discussed above for B7.

Complete CTL activation and effector function requires Th cell help through the interaction between the Th cell CD40L (CD40 ligand) molecule and the CD40 molecule expressed by DCs (Ridge et al., *Nature*, 393:474 (1998), Bennett et al., *Nature*, 393:478 (1998), Schoenberger et al., *Nature*, 393:480 (1998)). This mechanism of this costimulatory signal is likely to involve upregulation of B7 and associated IL-6/IL-12 production by the DC (APC). The CD40-CD40L interaction thus complements the signal 1 (antigen/MHC-TCR) and signal 2 (B7-CD28) interactions.

The use of anti-CD40 antibodies to stimulate DC cells directly, would be expected to

enhance a response to tumor antigens which are normally encountered outside of a inflammatory context or are presented by non-professional APCs (tumor cells). In these situations Th help and B7 costimulation signals are not provided. This mechanism might be used in the context of antigen pulsed DC based therapies or in situations where Th epitopes have not been defined within known TRA precursors.

A cancer associated antigen polypeptide, or a fragment thereof, also can be used to isolate their native binding partners. Isolation of such binding partners may be performed according to well-known methods. For example, isolated cancer associated antigen polypeptides can be attached to a substrate (e.g., chromatographic media, such as polystyrene beads, or a filter), and then a solution suspected of containing the binding partner may be applied to the substrate. If a binding partner which can interact with cancer associated antigen polypeptides is present in the solution, then it will bind to the substrate-bound cancer associated antigen polypeptide. The binding partner then may be isolated.

It will also be recognized that the invention embraces the use of the cancer associated antigen cDNA sequences in expression vectors, as well as to transfect host cells and cell lines, be these prokaryotic (e.g., *E. coli*), or eukaryotic (e.g., dendritic cells, B cells, CHO cells, COS cells, yeast expression systems and recombinant baculovirus expression in insect cells). Especially useful are mammalian cells such as human, mouse, hamster, pig, goat, primate, etc. They may be of a wide variety of tissue types, and include primary cells and cell lines. Specific examples include keratinocytes, peripheral blood leukocytes, bone marrow stem cells and embryonic stem cells. The expression vectors require that the pertinent sequence, i.e., those nucleic acids described *supra*, be operably linked to a promoter.

The invention also contemplates delivery of nucleic acids, polypeptides or peptides for vaccination. Delivery of polypeptides and peptides can be accomplished according to standard vaccination protocols which are well known in the art. In another embodiment, the delivery of nucleic acid is accomplished by *ex vivo* methods, i.e. by removing a cell from a subject, genetically engineering the cell to include a breast cancer associated antigen, and reintroducing the engineered cell into the subject. One example of such a procedure is outlined in U.S. Patent 5,399,346 and in exhibits submitted in the file history of that patent, all of which are publicly available documents. In general, it involves introduction *in vitro* of a functional copy of a gene into a cell(s) of a subject, and returning the genetically engineered cell(s) to the subject. The functional copy of the gene is under operable control of regulatory elements which permit expression of the gene in the genetically engineered cell(s). Numerous

transfection and transduction techniques as well as appropriate expression vectors are well known to those of ordinary skill in the art, some of which are described in PCT application WO95/00654. *In vivo* nucleic acid delivery using vectors such as viruses and targeted liposomes also is contemplated according to the invention.

5 In preferred embodiments, a virus vector for delivering a nucleic acid encoding a cancer associated antigen is selected from the group consisting of adenoviruses, adeno-associated viruses, poxviruses including vaccinia viruses and attenuated poxviruses, Semliki Forest virus, Venezuelan equine encephalitis virus, retroviruses, Sindbis virus, and Ty virus-like particle. Examples of viruses and virus-like particles which have been used to deliver  
10 exogenous nucleic acids include: replication-defective adenoviruses (e.g., Xiang et al., *Virology* 219:220-227, 1996; Eloit et al., *J. Virol.* 71:5375-5381, 1997; Chengalvala et al., *Vaccine* 15:335-339, 1997), a modified retrovirus (Townsend et al., *J. Virol.* 71:3365-3374, 1997), a nonreplicating retrovirus (Irwin et al., *J. Virol.* 68:5036-5044, 1994), a replication defective Semliki Forest virus (Zhao et al., *Proc. Natl. Acad. Sci. USA* 92:3009-3013, 1995),  
15 canarypox virus and highly attenuated vaccinia virus derivative (Paoletti, *Proc. Natl. Acad. Sci. USA* 93:11349-11353, 1996), non-replicative vaccinia virus (Moss, *Proc. Natl. Acad. Sci. USA* 93:11341-11348, 1996), replicative vaccinia virus (Moss, *Dev. Biol. Stand.* 82:55-63, 1994), Venezuelan equine encephalitis virus (Davis et al., *J. Virol.* 70:3781-3787, 1996), Sindbis virus (Pugachev et al., *Virology* 212:587-594, 1995), and Ty virus-like particle  
20 (Allsopp et al., *Eur. J. Immunol* 26:1951-1959, 1996). In preferred embodiments, the virus vector is an adenovirus.

Another preferred virus for certain applications is the adeno-associated virus, a double-stranded DNA virus. The adeno-associated virus is capable of infecting a wide range of cell types and species and can be engineered to be replication-deficient. It further has advantages,  
25 such as heat and lipid solvent stability, high transduction frequencies in cells of diverse lineages, including hematopoietic cells, and lack of superinfection inhibition thus allowing multiple series of transductions. The adeno-associated virus can integrate into human cellular DNA in a site-specific manner, thereby minimizing the possibility of insertional mutagenesis and variability of inserted gene expression. In addition, wild-type adeno-associated virus  
30 infections have been followed in tissue culture for greater than 100 passages in the absence of selective pressure, implying that the adeno-associated virus genomic integration is a relatively stable event. The adeno-associated virus can also function in an extrachromosomal fashion.

In general, other preferred viral vectors are based on non-cytopathic eukaryotic viruses

in which non-essential genes have been replaced with the gene of interest. Non-cytopathic viruses include retroviruses, the life cycle of which involves reverse transcription of genomic viral RNA into DNA with subsequent proviral integration into host cellular DNA.

Adenoviruses and retroviruses have been approved for human gene therapy trials. In general, the retroviruses are replication-deficient (i.e., capable of directing synthesis of the desired proteins, but incapable of manufacturing an infectious particle). Such genetically altered retroviral expression vectors have general utility for the high-efficiency transduction of genes *in vivo*. Standard protocols for producing replication-deficient retroviruses (including the steps of incorporation of exogenous genetic material into a plasmid, transfection of a packaging cell lined with plasmid, production of recombinant retroviruses by the packaging cell line, collection of viral particles from tissue culture media, and infection of the target cells with viral particles) are provided in Kriegler, M., "Gene Transfer and Expression, A Laboratory Manual," W.H. Freeman Co., New York (1990) and Murry, E.J. Ed. "Methods in Molecular Biology," vol. 7, Humana Press, Inc., Clifton, New Jersey (1991).

Preferably the foregoing nucleic acid delivery vectors: (1) contain exogenous genetic material that can be transcribed and translated in a mammalian cell and that can induce an immune response in a host, and (2) contain on a surface a ligand that selectively binds to a receptor on the surface of a target cell, such as a mammalian cell, and thereby gains entry to the target cell.

Various techniques may be employed for introducing nucleic acids of the invention into cells, depending on whether the nucleic acids are introduced *in vitro* or *in vivo* in a host. Such techniques include transfection of nucleic acid-CaPO<sub>4</sub> precipitates, transfection of nucleic acids associated with DEAE, transfection or infection with the foregoing viruses including the nucleic acid of interest, liposome mediated transfection, and the like. For certain uses, it is preferred to target the nucleic acid to particular cells. In such instances, a vehicle used for delivering a nucleic acid of the invention into a cell (e.g., a retrovirus, or other virus; a liposome) can have a targeting molecule attached thereto. For example, a molecule such as an antibody specific for a surface membrane protein on the target cell or a ligand for a receptor on the target cell can be bound to or incorporated within the nucleic acid delivery vehicle. Preferred antibodies include antibodies which selectively bind a cancer associated antigen, alone or as a complex with a MHC molecule. Especially preferred are monoclonal antibodies. Where liposomes are employed to deliver the nucleic acids of the invention, proteins which bind to a surface membrane protein associated with endocytosis

may be incorporated into the liposome formulation for targeting and/or to facilitate uptake. Such proteins include capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half life, and the like. Polymeric delivery systems also have been used successfully to deliver nucleic acids into cells, as is known by those skilled in the art. Such systems even permit oral delivery of nucleic acids.

When administered, the therapeutic compositions of the present invention can be administered in pharmaceutically acceptable preparations. Such preparations may routinely contain pharmaceutically acceptable concentrations of salt, buffering agents, preservatives, compatible carriers, supplementary immune potentiating agents such as adjuvants and cytokines and optionally other therapeutic agents.

The therapeutics of the invention can be administered by any conventional route, including injection or by gradual infusion over time. The administration may, for example, be oral, intravenous, intraperitoneal, intramuscular, intracavity, subcutaneous, or transdermal. When antibodies are used therapeutically, a preferred route of administration is by pulmonary aerosol. Techniques for preparing aerosol delivery systems containing antibodies are well known to those of skill in the art. Generally, such systems should utilize components which will not significantly impair the biological properties of the antibodies, such as the paratope binding capacity (see, for example, Sciarra and Cutie, "Aerosols," in Remington's Pharmaceutical Sciences, 18th edition, 1990, pp 1694-1712; incorporated by reference). Those of skill in the art can readily determine the various parameters and conditions for producing antibody aerosols without resort to undue experimentation. When using antisense preparations of the invention, slow intravenous administration is preferred.

The compositions of the invention are administered in effective amounts. An "effective amount" is that amount of a cancer associated antigen composition that alone, or together with further doses, produces the desired response, e.g. increases an immune response to the cancer associated antigen. In the case of treating a particular disease or condition characterized by expression of one or more cancer associated antigens, such as small cell lung cancer, the desired response is inhibiting the progression of the disease. This may involve only slowing the progression of the disease temporarily, although more preferably, it involves halting the progression of the disease permanently. This can be monitored by routine methods or can be monitored according to diagnostic methods of the invention discussed herein. The desired response to treatment of the disease or condition also can be delaying the onset or

even preventing the onset of the disease or condition.

Such amounts will depend, of course, on the particular condition being treated, the severity of the condition, the individual patient parameters including age, physical condition, size and weight, the duration of the treatment, the nature of concurrent therapy (if any), the specific route of administration and like factors within the knowledge and expertise of the health practitioner. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose of the individual components or combinations thereof be used, that is, the highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art, however, that a patient may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reasons.

The pharmaceutical compositions used in the foregoing methods preferably are sterile and contain an effective amount of cancer associated antigen or nucleic acid encoding cancer associated antigen for producing the desired response in a unit of weight or volume suitable for administration to a patient. The response can, for example, be measured by determining the immune response following administration of the cancer associated antigen composition via a reporter system by measuring downstream effects such as gene expression, or by measuring the physiological effects of the cancer associated antigen composition, such as regression of a tumor or decrease of disease symptoms. Other assays will be known to one of ordinary skill in the art and can be employed for measuring the level of the response.

The doses of cancer associated antigen compositions (e.g., polypeptide, peptide, antibody, cell or nucleic acid) administered to a subject can be chosen in accordance with different parameters, in particular in accordance with the mode of administration used and the state of the subject. Other factors include the desired period of treatment. In the event that a response in a subject is insufficient at the initial doses applied, higher doses (or effectively higher doses by a different, more localized delivery route) may be employed to the extent that patient tolerance permits.

In general, for treatments for eliciting or increasing an immune response, doses of cancer associated antigen are formulated and administered in doses between 1 ng and 1 mg, and preferably between 10 ng and 100 µg, according to any standard procedure in the art. Where nucleic acids encoding cancer associated antigen or variants thereof are employed, doses of between 1 ng and 0.1 mg generally will be formulated and administered according to standard procedures. Other protocols for the administration of cancer associated antigen

compositions will be known to one of ordinary skill in the art, in which the dose amount, schedule of injections, sites of injections, mode of administration (e.g., intra-tumoral) and the like vary from the foregoing. Administration of cancer associated antigen compositions to mammals other than humans, e.g. for testing purposes or veterinary therapeutic purposes, is carried out under substantially the same conditions as described above.

Where cancer associated antigen peptides are used for vaccination, modes of administration which effectively deliver the cancer associated antigen and adjuvant, such that an immune response to the antigen is increased, can be used. For administration of a cancer associated antigen peptide in adjuvant, preferred methods include intradermal, intravenous, intramuscular and subcutaneous administration. Although these are preferred embodiments, the invention is not limited by the particular modes of administration disclosed herein. Standard references in the art (e.g., *Remington's Pharmaceutical Sciences*, 18th edition, 1990) provide modes of administration and formulations for delivery of immunogens with adjuvant or in a non-adjuvant carrier.

When administered, the pharmaceutical preparations of the invention are applied in pharmaceutically-acceptable amounts and in pharmaceutically-acceptable compositions. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. Such preparations may routinely contain salts, buffering agents, preservatives, compatible carriers, and optionally other therapeutic agents. When used in medicine, the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically-acceptable salts thereof and are not excluded from the scope of the invention. Such pharmacologically and pharmaceutically-acceptable salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, maleic, acetic, salicylic, citric, formic, malonic, succinic, and the like. Also, pharmaceutically-acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts.

A small cell lung cancer associated antigen composition may be combined, if desired, with a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier" as used herein means one or more compatible solid or liquid fillers, diluents or encapsulating substances which are suitable for administration into a human. The term "carrier" denotes an organic or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions

also are capable of being co-mingled with the molecules of the present invention, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficacy.

The pharmaceutical compositions may contain suitable buffering agents, including:  
5 acetic acid in a salt; citric acid in a salt; boric acid in a salt; and phosphoric acid in a salt.

The pharmaceutical compositions also may contain, optionally, suitable preservatives, such as: benzalkonium chloride; chlorobutanol; parabens and thimerosal.

The pharmaceutical compositions may conveniently be presented in unit dosage form and may be prepared by any of the methods well-known in the art of pharmacy. All methods  
10 include the step of bringing the active agent into association with a carrier which constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product.

Compositions suitable for oral administration may be presented as discrete units, such  
15 as capsules, tablets, lozenges, each containing a predetermined amount of the active compound. Other compositions include suspensions in aqueous liquids or non-aqueous liquids such as a syrup, elixir or an emulsion.

Compositions suitable for parenteral administration conveniently comprise a sterile aqueous or non-aqueous preparation of breast cancer associated antigen polypeptides or  
20 nucleic acids, which is preferably isotonic with the blood of the recipient. This preparation may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation also may be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be  
25 employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono-or di-glycerides. In addition, fatty acids such as oleic acid may be used in the preparation of injectables. Carrier formulation suitable for oral, subcutaneous, intravenous, intramuscular, etc. administrations  
30 can be found in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA.

As used herein with respect to nucleic acids, the term "isolated" means: (i) amplified *in vitro* by, for example, polymerase chain reaction (PCR); (ii) recombinantly produced by cloning; (iii) purified, as by cleavage and gel separation; or (iv) synthesized by, for example,

chemical synthesis. An isolated nucleic acid is one which is readily manipulable by recombinant DNA techniques well known in the art. Thus, a nucleotide sequence contained in a vector in which 5' and 3' restriction sites are known or for which polymerase chain reaction (PCR) primer sequences have been disclosed is considered isolated but a nucleic acid sequence existing in its native state in its natural host is not. An isolated nucleic acid may be substantially purified, but need not be. For example, a nucleic acid that is isolated within a cloning or expression vector is not pure in that it may comprise only a tiny percentage of the material in the cell in which it resides. Such a nucleic acid is isolated, however, as the term is used herein because it is readily manipulable by standard techniques known to those of ordinary skill in the art. An isolated nucleic acid as used herein is not a naturally occurring chromosome.

As used herein with respect to polypeptides, "isolated" means separated from its native environment and present in sufficient quantity to permit its identification or use. Isolated, when referring to a protein or polypeptide, means, for example: (i) selectively produced by expression cloning or (ii) purified as by chromatography or electrophoresis. Isolated proteins or polypeptides may, but need not be, substantially pure. The term "substantially pure" means that the proteins or polypeptides are essentially free of other substances with which they may be found in nature or *in vivo* systems to an extent practical and appropriate for their intended use. Substantially pure polypeptides may be produced by techniques well known in the art. Because an isolated protein may be admixed with a pharmaceutically acceptable carrier in a pharmaceutical preparation, the protein may comprise only a small percentage by weight of the preparation. The protein is nonetheless isolated in that it has been separated from the substances with which it may be associated in living systems, i.e. isolated from other proteins.

## Examples

### Methods and Materials

#### *Cell lines, tissues, and patient sera*

Cell lines were obtained from the repository maintained at the Ludwig Institute for Cancer Research (LICR), New York Branch at the Memorial Sloan-Kettering Cancer Center (MSKCC), or obtained from American Tissue Culture Collection. Eleven SCLC cell lines were used including 9 classical (SK-LC-13, NCI-H69, -H128, -H146, -H187, -H209, -H378, -H889, -H740) and 2 variant (NCI-H82, -H526) forms. The variant SCLC lines differ from the classical lines in lacking or having diminished neuroendocrine features and with regard to

other biochemical, morphological and growth properties (Carney et al., *Cancer Res.* 45:2913-2923, 1985; Park et al., *Cancer Res.* 47:6710-6718, 1987). Normal and tumor tissues were obtained from the departments of Pathology in the New York Presbyterian Hospital (NYPH) and the MSKCC. Patient sera were obtained from the Department of Medicine, NYPH, and  
5 from the LICR Melbourne Branch, Australia.

### *Immunoscreening of the SCLC cell line libraries and characterization of immunoreactive clones*

Construction of cDNA expression libraries from the NCI-H740 and SK-LC-13 SCLC  
10 cell lines in the  $\lambda$ -ZAP vector (Stratagene) and immunoscreening were done as previously described (Old and Chen, *J. Exp. Med.*, 187:1163-7, 1998; Chen et al., *Proc. Nat'l. Acad. Sci. U S A*, 95: 6919-23, 1998), with the following modifications. Sera from five SCLC patients (Lu94, Lu100, Lu101, Lu104, Lu113) were pooled and absorbed as previously described Scanlan et al., *Int. J. Cancer* 76:652-658, 1998). The pooled serum was diluted 1:200 (final  
15 dilution 1:1000 for each serum) in TBS containing 1% BSA and 0.02% NaN<sub>3</sub> and was used to screen  $5.6 \times 10^5$  pfu of the NCI-H740 library. The same serum was used for the SK-LC-13 library of which  $2.2 \times 10^5$  pfu was screened. Immunoreactive clones were isolated and sequence analyzed as previously described (Chen et al., 1998). Selected immunoreactive clones were subsequently tested for reactivity against sera at various dilutions from individual  
20 lung cancer patients and normals using the same plaque assay. A  $\lambda$ -ZAP clone without an insert was co-plated and included in the screen as a negative control.

### *RT-PCR analysis*

Reverse transcription was performed with total RNA isolated from tissue or cell lines  
25 by the Guanidium thiocyanate / CsCl method. Primers used to amplify *ZIC2* were designed based on the published sequence (AF104902) and our results. ZIC2A1: 5'-CATGAATATGAACATGGGTATGAACATGG (SEQ ID NO:1); ZIC2B1: 5'-TCGCAGCCCTCAAACCTCACACTG (SEQ ID NO:2). Conditions for amplification were as follows: Initial denaturation and AmpliTaq Gold (Perkin Elmer) activation; 94°C, 10',  
30 Amplification: 94°C, 1'; 60°C, 1'; 72°C, 1'; for 35 cycles, followed by a 6', 72°C incubation. Amplification products were analyzed by agarose gel electrophoresis and visualized by EtBr staining.

### *Northern blot analysis*

Adult normal tissue mRNA blots were obtained from Clontech, Inc. and contained 2 g polyA<sup>+</sup> RNA per lane. Lung cancer cell line total RNA was isolated as described above and polyA<sup>+</sup> mRNA was prepared using the Microfast Track kit (Invitrogen). Two grams of mRNA or 10 g of total RNA was transferred to nylon membranes (Schleicher and Schuell) following denaturing gel electrophoresis. Hybridizations and washes were carried out under high stringency conditions in ExpressHyb buffer (Clontech) using hybridization and washing conditions described by the manufacturer. The probes used for northern blot analysis were the following. *SOX2*: 450 bp fragment (nucleotides 630-1080); *SOX1*: 751 bp fragment (nucleotides 1520-2271); *SOX3*: 330 bp fragment (nucleotides 442-772); *SOX21*: 680 bp fragment (nucleotides 2720-3400); and *ID4*: full-length cDNA (1322bp).

### **Example 1: Isolation of Immunoreactive clones from SCLC cell lines by SEREX**

SEREX analysis of the SCLC cell line NCI-H740 with a pool of five sera from SCLC patients at 1:10<sup>3</sup> dilution resulted in the isolation of 37 clones coding for 8 known gene products (Table 1a). These eight genes were given SEREX gene designations of *NY-SCLC-1* to *NY-SCLC-8*.

**Table 1a. Genes isolated by SEREX analysis of the small cell lung cancer cell line NCI-H740**

SEQ ID NO:	Gene Designation	Gene/Sequence Identity [GenBank Accession No.]	Number of clones (% of total)
3	<i>NY-SCLC-1</i>	<i>SOX2</i> [Z31560]	19 (51%)
4	<i>NY-SCLC-2</i>	<i>SOX1</i> [Y13436]	1 (3%)
5	<i>NY-SCLC-3</i>	<i>ZIC2</i> [AF104902]	9 (24%)
6	<i>NY-SCLC-4</i>	<i>ID4</i> [U28368]	2 (5%)
7	<i>NY-SCLC-5</i>	<i>MAZ</i> [M94046]	1 (3%)
8	<i>NY-SCLC-6</i>	<i>MPP11</i> [X98260]	3 (8%)
9	<i>NY-SCLC-7</i>	<i>eIF2B</i> [U23028]	1 (3%)
10	<i>NY-SCLC-8</i>	<i>RBP-1</i> [L07872]	1 (3%)
			Total: 37

The most frequently isolated genes were *SOX2* and *ZIC2*, comprising 51% and 24% of all clones. A single clone corresponding to *SOX1* was also isolated from this library. *SOX-*

and *ZIC2*-encoding clones showed very strong immunoreactivity with the SCLC patient sera. Other genes isolated included *ID4*, *MPP11*, *MAZ*, *eIF2B* and *RBP-1*. *ID4* protein is a member of the dominant negative helix-loop-helix (HLH) proteins. This protein can interact with other HLH proteins such as the one encoded by *Archaeo-Scute* and by virtue of not containing a DNA binding domain it acts as a repressor (Riechmann, et al., *Nucleic Acids Res.*, 22: 749-55, 1994). The mRNA expression pattern of *ID4* in normal tissues was found to be universal by Northern blot analysis. Seroreactivity against *ID4* was moderate at 1:10<sup>3</sup> sera dilution. *MPP11* is another HLH protein-binding factor, and it has also been isolated from HeLa cells by M-phase protein-recognizing antibodies (Shoji, et al., *J. Biol. Chem.*, 270:24818-25, 1995; Matsumoto-Taniura, et al. *Mol. Biol. Cell*, 7: 1455-69, 1996). Seroreactivity against *MPP11* was strong at a 1:1000 dilution of the SCLC sera. This antigen was also identified by SEREX analysis of gastric and breast cancer and is universally expressed. Other genes isolated from NCI-H740 -- the myc-associated Zinc-finger protein *MAZ*, the eukaryotic translation initiation factor *eIF2B* and the J- $\kappa$  recombination signal binding protein (*RBP-1*) -- were also previously identified by SEREX. *MAZ*, *eIF2B* and *RBP-1* are expressed in multiple normal adult tissues.

The SEREX analysis of the second SCLC line SK-LC-13 with the same pooled sera from SCLC patients resulted in the identification of 14 clones corresponding to 10 genes (Table 1b), 4 of which were identical to those isolated from NCI-H740 and 6 were distinct (NY-SCLC-9 to NY-SCLC-14).

**Table 1b.** Genes isolated by SEREX analysis of the small cell lung cancer cell line SK-LC-13

SEQ ID NO:	Gene Designation	Gene/Sequence Identity [GenBank Accession No.]	Number of clones (% of total)
3	NY-SCLC-1	SOX2 [Z31560]	2 (14%)
11	NY-SCLC-9	SOX3 [X71135]	1 (7%)
12	NY-SCLC-10	SOX21 [AF107044]	1 (7%)
5	NY-SCLC-3	ZIC2 [AF104902]	2 (14%)
6	NY-SCLC-4	ID4 [U28368]	1 (7%)
8	NY-SCLC-6	MPP11 [X98260]	3 (21%)
13	NY-SCLC-11	KIAA0963 [AB023180.1]	1 (7%)
14	NY-SCLC-12	LAG-3 [X51985]	1 (7%)
15, 16	NY-SCLC-13	DKFZp434C196	1 (7%)

		[AL133561.1]	
17	<i>NY-SCLC-14</i>	<i>Novel-2</i>	1 (7%)
			Total: 14

*SOX2* was isolated twice and in addition *SOX3* and *SOX21* were isolated, each represented by a single clone. *ZIC2* was isolated twice. Other genes isolated that were identical to those from the NCI-H740 library included *ID4*, isolated once, and *MPP11*, which was represented by three immunoreactive clones. Among other genes identified, *NY-SCLC-11* (*KIAA0963*) is an unknown gene with identical EST sequences derived from many tissues. Two novel genes (*NY-SCLC-13*) and (*NY-SCLC-14*) were isolated, one of which (*NY-SCLC-14*) showed no sequence identity to current GenBank entries. These two genes were intriguing in that their DNA sequences contain homopolymers of 24 bp and 6 bp repeats and would encode tandem octapeptides and dipeptides, respectively. *NY-SCLC-12*, lymphocyte activation gene-3 (*LAG-3*), is related to CD4 and has a restricted tissue expression pattern, possibly representing a differentiation antigen of lymphoid origin (Triebel, et al., *J. Exp. Med.*, 171: 1393-1405, 1990).

## **Example 2: Immunodominant epitopes of ZIC2 and the SOX proteins**

Of 11 *ZIC2* clones isolated, 7 clones were sequenced and 4 were evaluated by restriction mapping. The longest *ZIC2* clone (NCI-H740 #32) was ~2.6kb, the sequence of which extends beyond both 5' and 3' sequences of the *ZIC2* cDNA entry in the GenBank (AF104902). The shortest clone (NCI-H740 #41) migrated as a ~1kb band on agarose gels and its 5' end corresponded to nucleotide position 692 (amino acid residue 231) of AF104902. Reactivity of this clone with SCLC sera was comparable to other larger clones. As the intensity of the reactivity of this shorter clone was comparable to that of other larger *ZIC2* clones, the seroreactive epitope(s) of *ZIC2* polypeptide (SEQ ID NO:22) reside between amino acid residue 231 and the C-terminal end (amino acid residue 533).

Of the 24 *SOX* genes, 8 *SOX2* clones and the *SOX1*, *SOX3* and *SOX21* clones were sequence analyzed while the remaining 13 *SOX2* clones were analyzed and confirmed by restriction mapping. All *SOX2* clones contained the full size cDNA (1085bp) and the longest clone (NCI-H740 #2) had 54 additional nucleotides at its 5' untranslated region as compared to the *SOX2* GenBank entry (Accession Number Z31560). The two *SOX1* and *SOX3* clones contained truncated cDNA inserts which lacked sequences 5' to those encoding the HMG-box

while the *SOX21* clone encoded the full length SOX21 protein, which has only 5 residues N-terminal to its HMG-box (Fig. 1). The most conserved region among these *SOX* cDNA clones is thus the HMG-box-encoding region which is 88 to 96% identical among the SOX Group B proteins. All sera that reacted with SOX1 also reacted with SOX2, SOX3 and SOX21 (see below), suggesting that at least part of the immunoreactivity of SCLC patient sera is directed against the conserved HMG-box of the SOX proteins.

### **Example 3: *ZIC2* is expressed exclusively in brain, testis and tumors**

*ZIC2* gene expression was analyzed by RT-PCR. The RNA quality was confirmed by successful amplification of *p53* exons 5 and 6. Among normal tissues *ZIC2* mRNA was only detectable in brain and to a lesser extent in testis but not in skin, kidney, small intestine, pancreas, uterus and lung. Of 11 SCLC cell lines analyzed, all 9 classical SCLC lines (SK-LC-13, NCI-H69, -H128, -H146, -H187, -H209, -H378, -H889, -H740) had detectable *ZIC2* mRNA while two variant SCLC cell lines (NCI-H82 and NCI-H526) showed no or minimal expression. Among other cell lines, *ZIC2* mRNA could be amplified in 100% (7/7) of non-small cell lung tumor cell lines and 83% (10/12) of melanoma cell lines (Table 2). Among tumor tissues, 50% (5/10) of melanoma, 50% (2/4) of colon cancer, 75% (3/4) of breast cancer, 86% (12/14) of head and neck cancer, 66% (6/9) of lung cancer, 50% (7/14) of transitional cancer, 50% (1/2) of leiomyosarcoma and 100% (2/2) of synovial sarcoma samples had detectable *ZIC2* mRNA (**Table 2**).

**Table 2. *ZIC2* gene expression in cancer cell lines and tumor samples**

<b>TUMOR CELL LINE</b>	<b><i>ZIC2</i> mRNA EXPRESSION</b>
Melanoma	10/12 (83%)
NSCLC	7/7 (100%)

  

<b>TUMOR TYPE</b>	<b><i>ZIC2</i> mRNA EXPRESSION</b>
Melanoma	5/10 (50%)
Colon cancer	2/4 (50%)
Breast cancer	3/4 (75%)
Head & neck cancer	12/14 (86%)
Lung cancer	6/9 (66%)
Transitional cancer	7/14 (50%)

Leiomyosarcoma	1/2 (50%)
Synovial sarcoma	2/2 (100%)

#### **Example 4: SOX gene expression characteristics**

Since *SOX Group B* genes are intronless, RT-PCR results using tissue RNA were often difficult to interpret due to the genomic DNA contamination of RNA samples. Therefore, their gene expression was evaluated by Northern blot analysis. An  $\alpha$ -actin probe was used to confirm the RNA quality and quantity. Northern blots were exposed for 24 h (*SOX2* - SCLC blot), 72 h (*SOX1*), or 1 week (*SOX3*, *SOX21* and *SOX2* - normal tissue blot).

Among normal tissues *SOX2* mRNA could be detected in brain, testis and prostate, and at lower levels in small intestine and colon but not in heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, ovary and peripheral blood leukocytes. *SOX1*, *SOX3* and *SOX21* mRNA were not detected in normal adult tissues, which is consistent with the current literature. *SOX Group B* expression in tumor cell lines was also examined. *SOX2* was expressed in 5 of 10 SCLC cell lines (NCI-H69, NCI-H146, NCI-H378, NCI-H740 and SK-LC-13). *SOX2* message was not detected in the three non-SCLC cell lines SK-LC-7, 8 and 17 or in the 8 melanoma cell lines SK-MEL-10, 12, 14, 24, 26, 28, 37 and Mz19. *SOX1* mRNA was detected in 4 of 10 SCLC cell lines (NCI-H187, NCI-H209, NCI-H378 and SK-LC-13) while *SOX3* mRNA could be detected in 2 of 10 SCLC cell lines (NCI-H740, and as a weaker signal in SK-LC-13). *SOX1* and *SOX3* required longer exposure times than *SOX2*, indicating their expression levels are lower than that of *SOX2*. *SOX21* mRNA was not detected after prolonged exposure (1 week), indicating no or low levels of expression. Two variant SCLC cell lines, NCI-H82 and NCI-H526, had no detectable *SOX Group B* expression.

#### **Example 5: SCLC patient sera contain high-titer antibodies to SOX and ZIC2 proteins**

Reactivity to phage clones containing *SOX1*, 2, 3, 21 and *ZIC2* was titrated against 17 SCLC patient sera and 16 normal adult sera. ZAP phages with no insert were mixed with the test clone and served as internal negative controls, visible as a background at 1:10<sup>4</sup> serodilution on Lu113. Assays were scored positive only when test clones could be clearly distinguished from the control phages.

Only one of the 16 normal sera showed weak reactivity against *SOX2* at a titer of 1:1000. In contrast, 7 of 17 patients (41%) had antibodies reactive with *SOX1* and *SOX2*

containing phagemids while 29% (5/17) and 35% (6/17) had antibodies to SOX3 and SOX21 respectively. 29% (5/17) of patients had detectable anti-ZIC2 antibodies. The antibody titers measured up to  $1:10^6$  (Table 3). All five patient sera that had antibodies against ZIC2 also reacted with SOX proteins at varying titers; one (Lu113) was reactive at  $1:10^6$  while another (Lu139) was reactive only at a  $1:10^3$  dilution. Two patients (Lu100 and A6) had antibodies against SOX1 and SOX2 proteins at  $1:10^5$  but no antibodies against ZIC2 even at  $1:10^3$  dilution (Table 3).

**Table 3.** SOX and ZIC2 Reactivity of Small Cell Lung Cancer Patient Sera

	Protein: Serum:	SOX1	SOX2	SOX3	SOX21	ZIC2
1	Lu 94*	$1:10^5$	$1:10^5$	$1:10^4$	$1:10^4$	$1:10^5$
2	Lu 100*	$1:10^5$	$1:10^5$	$1:10^4$	$1:10^4$	-
3	Lu 101*	-	-	-	-	-
4	Lu 104*	-	-	-	-	-
5	Lu 113*	$1:10^6$	$1:10^6$	$1:10^5$	$1:10^5$	$1:10^6$
6	Lu 139	$1:10^3$	$1:10^3$	-	-	$1:10^6$
7	Lu 159	-	-	-	-	-
8	A1	$1:10^5$	$1:10^6$	$1:10^5$	$1:10^5$	$1:10^4$
9	A2	-	-	-	-	-
10	A3	-	-	-	-	-
11	A4	-	-	-	-	-
12	A5	-	-	-	-	-
13	A6	$1:10^5$	$1:10^5$	$1:10^3$	$1:10^4$	-
14	A7	-	-	-	-	-
15	A8	-	-	-	-	-
16	A9	-	-	-	-	-
17	A10	$1:10^4$	$1:10^4$	-	$1:10^3$	$1:10^4$
		7/17(41%)	7/17(41%)	5/17(29%)	6/17(35%)	5/17(29%)

\*Pooled sera used for SEREX analysis of the SCLC cell lines

All patients who had antibodies against SOX3 or SOX21 had antibodies at higher titers against SOX1 and SOX2. The presence of consistently higher titer antibodies against SOX1 and SOX2 suggests SOX1 and/or 2 as the main immunogenic tumor antigen in these

patients, whereas the seroreactivity to SOX3 and SOX21 might be secondary to the share antigenic epitopes located within the highly conserved HMG-box among SOX proteins.

From the immunological standpoint, the high frequency and high titers of anti-SOX and anti-ZIC2 antibodies in these SCLC patients is striking. Anti-ZIC2 antibody was  
5 observed in 29% and anti-SOX antibody was observed in 41% of the SCLC sera tested. These sera were collected from a heterogeneous group of SCLC patients who were at different stages of their diseases, receiving various treatments, and with variable responses; one of the antibody-positive patients (Lu113) had no clinical evidence of residual disease when serum was collected and had subsequent recurrence of tumor. This means that if serum is collected  
10 from untreated cases of SCLC, the frequency of detecting anti-SOX and anti-ZIC2 antibodies can be substantially higher than the 30-40% rate found in this study. This frequency is significantly higher than the antibody responses seen against most other SEREX-defined antigens. Scanlan et al. (*Int. J. Cancer* 76:652-658, 1998) have evaluated large panels of SEREX antigens for seroreactivity in cancer and normal patients. It was found that antigens  
15 that elicit cancer-specific antibody responses tend to have detectable seral antibody in up to 20-25% of tumor patients, rarely exceeding 25%. In this regard, the immunogenicity of SOX and ZIC2 antigens in these patients are exceptional and this indicates that an antibody-based assay can be useful in the diagnosis of SCLC, e.g. as a screening test for the high-risk group. Also, for SCLC cases that have been shown to have high-titer antibodies, the titer of the  
20 antibody can be correlated to the clinical progression/remission of the disease. If the presence of antibody is dependent on the tumor load, as has been shown for another SEREX-defined antigen, NY-ESO-1 (Stockert et al., *J. Exp. Med.* 187:1349-1354, 1998), antibody monitoring in these patients may also be of clinical value.

In addition to its immunodiagnostic potential, *SOX group B* and *ZIC2* products can be  
25 used as targets for cancer vaccines. The expression of these genes in brain may be a concern, particularly given the clinically-recognized paraneoplastic syndromes and their correlation to the aberrant expression of neural antigens in SCLC (Dalmau & Posner, *Arch. Neurol.* 56:405-408, 1999; Posner & Dalmau, *Curr. Opin. Immunol.* 9:723-729, 1997). However, despite the presence of high-titer anti-SOX and anti-ZIC2 antibodies, none of the seven antibody-positive  
30 patients in this study had neurological manifestations of the disease. In fact, the only patient in this study with paraneoplastic disease involving the cerebellum (patient A9) had no detectable anti-SOX Group B or anti-ZIC2 antibodies. The immune responses toward these antigens thus may not lead to autoimmune neurological disorders in most patients. Since *SOX*

and *ZIC2* genes are conserved in mice, preclinical studies can be carried out by SOX and/or *ZIC2* vaccination in these experimental models. Indeed, HuD antigen, one of the antigens associated with paraneoplastic syndromes, has recently been used as a vaccine target in the murine model of small cell lung cancer, and antitumor activity was observed without neurological disease (Carpentier et al., *Clin. Cancer Res.* 4:2819-2824, 1998; Ohwada et al., *Am. J. Respir. Cell. Mol. Biol.* 21:37-43, 1999).

#### **Example 6: Preparation of recombinant cancer associated antigens**

To facilitate screening of patients' sera for antibodies reactive with cancer associated antigens, for example by ELISA, recombinant proteins are prepared according to standard procedures. In one method, the clones encoding cancer associated antigens are subcloned into a baculovirus expression vector, and the recombinant expression vectors are introduced into appropriate insect cells. Baculovirus/insect cloning systems are preferred because post-translational modifications are carried out in the insect cells. Another preferred eukaryotic system is the *Drosophila* Expression System from Invitrogen. Clones which express high amounts of the recombinant protein are selected and used to produce the recombinant proteins. The recombinant proteins are tested for antibody recognition using serum from the patient which was used to isolate the particular clone, or in the case of cancer associated antigens recognized by allogeneic sera, by the sera from any of the patients used to isolate the clones or sera which recognize the clones' gene products.

Alternatively, the cancer associated antigen clones are inserted into a prokaryotic expression vector for production of recombinant proteins in bacteria. Other systems, including yeast expression systems and mammalian cell culture systems also can be used.

#### **Example 7: Preparation of antibodies to cancer associated antigens**

The recombinant cancer associated antigens produced as in Example 6 above are used to generate polyclonal antisera and monoclonal antibodies according to standard procedures. The antisera and antibodies so produced are tested for correct recognition of the cancer associated antigens by using the antisera/antibodies in assays of cell extracts of patients known to express the particular cancer associated antigen (e.g. an ELISA assay). These antibodies can be used for experimental purposes (e.g. localization of the cancer associated antigens, immunoprecipitations, Western blots, etc.) as well as diagnostic purposes (e.g., testing extracts of tissue biopsies, testing for the presence of cancer associated antigens).

The antibodies are useful for accurate and simple typing of small cell lung cancer tissue samples for expression of *SOX Group B* and *ZIC2* genes. SCLC is usually diagnosed by endoscopic biopsies rather than surgical resection, and an adequate specimen for RNA extraction and RT-PCR typing may not be obtained in every case. These difficulties are further complicated by the fact that *SOX Group B* genes are intronless, and RT-PCR is often unreliable. The best technique to type the expression of these genes and circumvent these problems is by immunohistochemical analysis with specific antibody reagents.

**Example 8: Expression of cancer associated antigens in cancers of similar and different origin.**

The expression of one or more of the cancer associated antigens is tested in a range of tumor samples to determine which, if any, other malignancies should be diagnosed and/or treated by the methods described herein. Tumor cell lines and tumor samples are tested for cancer associated antigen expression, preferably by RT-PCR according to standard procedures, e.g., as described for *ZIC2* expression in Example 3 above. Northern blots also are used to test the expression of the cancer associated antigens. Antibody based assays, such as ELISA and western blot, also can be used to determine protein expression. A preferred method of testing expression of cancer associated antigens (in other cancers and in additional same type cancer patients) is allogeneic serotyping using a modified SEREX protocol (as described above).

In all of the foregoing, extracts from the tumors of patients who provided sera for the initial isolation of the cancer associated antigens are used as positive controls. The cells containing recombinant expression vectors described in the Examples above also can be used as positive controls.

The results generated from the foregoing experiments provide panels of multiple cancer associated nucleic acids and/or polypeptides for use in diagnostic (e.g. determining the existence of cancer, determining the prognosis of a patient undergoing therapy, etc.) and therapeutic methods (e.g., vaccine composition, etc.).

**Example 9: HLA typing of patients positive for cancer associated antigens**

To determine which HLA molecules present peptides derived from the cancer associated antigens of the invention, cells of the patients which express the cancer associated antigens are HLA typed. Peripheral blood lymphocytes are taken from the patient and typed

for HLA class I or class II, as well as for the particular subtype of class I or class II. Tumor biopsy samples also can be used for typing. HLA typing can be carried out by any of the standard methods in the art of clinical immunology, such as by recognition by specific monoclonal antibodies, or by HLA allele-specific PCR (e.g. as described in WO97/31126).

5

**Example 10: Characterization of cancer associated antigen peptides presented by MHC class I and class II molecules.**

Antigens which provoke an antibody response in a subject may also provoke a cell-mediated immune response. Cells process proteins into peptides for presentation on MHC class I or class II molecules on the cell surface for immune surveillance. Peptides presented by certain MHC/HLA molecules generally conform to motifs. These motifs are known in some cases, and can be used to screen the small cell lung cancer associated antigens for the presence of potential class I and/or class II peptides. Summaries of class I and class II motifs have been published (e.g., Rammensee et al., *Immunogenetics* 41:178-228, 1995). Based on the results of experiments such as those described above, the HLA types which present the individual breast cancer associated antigens are known. Motifs of peptides presented by these HLA molecules thus are preferentially searched.

One also can search for class I and class II motifs using computer algorithms. For example, computer programs for predicting potential CTL epitopes based on known class I motifs has been described (*see, e.g.*, Parker et al, *J. Immunol.* 152:163, 1994; D'Amato et al., *Human Immunol.* 43:13-18, 1995; Drijfhout et al., *Human Immunol.* 43:1-12, 1995). Computer programs for predicting potential T cell epitopes based on known class II motifs has also been described (*see, e.g.* Sturniolo et al., *Nat Biotechnol* 17(6):555-61, 1999). HLA binding predictions can conveniently be made using an algorithm available via the Internet on the National Institutes of Health World Wide Web site at URL <http://bimas.dcrt.nih.gov> . See also the website of: SYFPEITHI: An Internet Database for MHC Ligands and Peptide Motifs (access via <http://www.uni-tuebingen.de/uni/kxi/> or <http://134.2.96.221/scripts/hlaserver.dll/EpPredict.htm>. Methods for determining HLA class II peptides and making substitutions thereto are also known (e.g. Strominger and Wucherpennig (PCT/US96/03182)).

**Example 11: Identification of the portion of a cancer associated polypeptide encoding an antigen**

To determine if the cancer associated antigens isolated as described above can provoke a cytolytic T lymphocyte response, the following method is performed. CTL clones are generated by stimulating the peripheral blood lymphocytes (PBLs) of a patient with autologous normal cells transfected with one of the clones encoding a cancer associated antigen polypeptide or with irradiated PBLs loaded with synthetic peptides corresponding to the putative protein and matching the consensus for the appropriate HLA class I molecule (as described above) to localize an antigenic peptide within the cancer associated antigen clone (see, e.g., Knuth et al., *Proc. Natl. Acad. Sci. USA* 81:3511-3515, 1984; van der Bruggen et al., *Eur. J. Immunol.* 24:3038-3043, 1994). These CTL clones are screened for specificity against COS cells transfected with the cancer associated antigen clone and autologous HLA alleles as described by Brichard et al. (*Eur. J. Immunol.* 26:224-230, 1996). CTL recognition of a cancer associated antigen is determined by measuring release of TNF from the cytolytic T lymphocyte or by  $^{51}\text{Cr}$  release assay (Herin et al., *Int. J. Cancer* 39:390-396, 1987). If a CTL clone specifically recognizes a transfected COS cell, then shorter fragments of the cancer associated antigen clone transfected in that COS cell are tested to identify the region of the gene that encodes the peptide. Fragments of the cancer associated antigen clone are prepared by exonuclease III digestion or other standard molecular biology methods. Synthetic peptides are prepared to confirm the exact sequence of the antigen.

Optionally, shorter fragments of cancer associated antigen cDNAs are generated by PCR. Shorter fragments are used to provoke TNF release or  $^{51}\text{Cr}$  release as above.

Synthetic peptides corresponding to portions of the shortest fragment of the cancer associated antigen clone which provokes TNF release are prepared. Progressively shorter peptides are synthesized to determine the optimal cancer associated antigen tumor rejection antigen peptides for a given HLA molecule.

A similar method is performed to determine if the cancer associated antigen contains one or more HLA class II peptides recognized by T cells. One can search the sequence of the cancer associated antigen polypeptides for HLA class II motifs as described above. In contrast to class I peptides, class II peptides are presented by a limited number of cell types. Thus for these experiments, dendritic cells or B cell clones which express HLA class II molecules preferably are used.

#### **Table 4: Sequence homologies**

SEQ ID NO:15 (NY-SCLC-13 5' SEQUENCE)

AL133561.1, AC007324.53, AP000552.1, AP000550.1, AC007708.13, AC009288.12, AC007325.49, AC008103.23, AC008079.22, AC008018.18, AC007731.11, AC005500, AC012398.3, AC008132.33, AC011718.2, AL117481.1, AE001958.1, AJ243721.1, X70255, X54676, AL110383.1, AL041090.1, 5 AW261390.1, AI904151.1, AA314127, H29680, H08571, R60682, R54134, R50027, R19696, R18168, R12223, F13183, F12174, F07553, F07164, F05322, F05321, F05267, F05235, T33549, Z43231, AI828436.1, AW226624.1, AW012831.1, AW012161.1, AI874452.1, AI391139, AI225578, AI099322, AA510280, AA475860, AA276058, AA277960, AA239475, AA139948, AA106968, AA073333, AA066928, AA002337, W18896, W07975, AW148528.1, AI934011.1, AI885936.1, AI885982.1, AI824746.1, AI801523.1, 10 AI741661.1, AI679504.1, AI589998.1, AI567632.1, AI564170.1, AI520793.1, AA677535, AA292543, F06393, AW142285.1, AW140928.1, AA520277.

SEQ ID NO:16 (NY-SCLC-13 3' SEQUENCE)

15 AL133561.1, AC007324.53, AP000552.1, AP000550.1, AC007708.13, AC009288.12, AC007325.49, AC008103.23, AC008079.22, AC008018.18, AC005500, AC007731.11, AC012398.3, AC008132.33, AC011718.2, AL117481.1, AE001958.1, X70255, X54676, AF022185, U00016, AL110383.1, AL041090.1, AW261390.1, AI904151.1, AW012161.1, AI391139, AI741661.1, AW142285.1, AW140928.1, AA520277.

20 SEQ ID NO:17 (NY-SCLC-14)

X14112, D10879, Z68873.1, AJ009970.1, AF077000, M11043, AC004093, L04961, AC008124.8, AC005742, AC000395, AL023802.1, U44088, AL031258.8, U92983, Z50194, Z63758, M55701, M80829, AF192802.1, Z84494.1, AC005387, AC004490, Z93784.1, AC003976, M69157, AL031864.1, M11041, AF131866.1, 25 AL023284.1, AF039833, U62317, NM\_003980.1, AF132809.1, NM\_003632.1, U38195, U38193, S44199, AB000634, NM\_003459.1, NM\_006245.1, D78360, AC004471, U04357, L77570, U52112, M97881, L22206, NM\_004565.1, AB018269.1, AE001198, AF022844, Z82173.2, AF167560.1, AC007032.2, AB020714.1, AF037372, AC002984, U81524, U63850, Z64726, X80330, AL110210.1, AL096857.1, AL031597.7, AL021579.1, AC005932, M63138, M28265, X80327, L14589, AC011718.2, Z92546.2, AC008018.18, 30 AP000353.1, AC004148, Y08701, AF023268, U77716, U46921, U46920, AC006549.27, Z99757.12, AC005817.6, AL035090.10, AC003063.7, AC004828.2, AC006547.9, AC000097, AF051345, Z94162.1, U34879, M84472.1, AF190826.1, M73779, AC002094, AW001248.1, AI863828.1, AI858055.1, AI813670.1, AI684429.1, AI277482, AI580934.1, AA472637, W64993, AW043820.1, AW028151.1, AI949719.1, AI887909.1, AI805058.1, AI804955.1, AI798900.1, AI741492.1, AI582191.1, AI348656, AI336325, 35 AI299745, AI276119, AI269740, AI262960, AI200633, AI097473, AA884197, AA527274, AA480684, W68353, AI931453.1, AA726490, W98413, AW263065.1, AW211900.1, AI006238, AA255056, AA238335, H27099, AA673074, AW139762.1, AL047473.1, AW223562.1, AW066814.1, AW031777.1, AI782249.1, AI774556.1, AI586471.1, AA139570, AI923922.1, AV390350.1, AA505122, AA380178, AI853595.1, AI851994.1, AI846520.1, AI154485, AI007056, AA467529, AA274838, AA261057, AA032648, W70846, 40 W71079, AA323008, C95416.1, AW210204.1, AA718506, AL042695.1, T25132, AI997515.1, AW205598.1, AI686223.1, AI590082.1, AI378378, AI318623, AI318236, AI201238, AI200900, AI190426, AI022738, AA916388, AA865035, AA845480, AA778028, AA744509, AA679215, AA558436, AA456062, AA418017, AA328237, AA159291, AA129371, N33970, H43255, T77577, AI890886.1, AA292501, AI379199.1, AI831459.1.

45

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

50 All references disclosed herein are incorporated by reference in their entirety.

We claim:

**Claims**

1. A method of diagnosing a disorder characterized by expression of a human cancer associated antigen precursor coded for by a nucleic acid molecule, comprising:

contacting a biological sample isolated from a subject with an agent that specifically  
5 binds to the nucleic acid molecule, an expression product thereof, or a fragment of an expression product thereof complexed with an HLA molecule, wherein the nucleic acid molecule is a NA Group 1 nucleic acid molecule, and

determining the interaction between the agent and the nucleic acid molecule or the expression product as a determination of the disorder.

10

2. The method of claim 1, wherein the agent is selected from the group consisting of

(a) a nucleic acid molecule comprising NA group 1 nucleic acid molecules or a fragment thereof,

(b) a nucleic acid molecule comprising NA group 3 nucleic acid molecules or a  
15 fragment thereof,

(c) a nucleic acid molecule comprising NA group 5 nucleic acid molecules or a fragment thereof,

(d) an antibody that binds to an expression product of NA group 1 nucleic acids,

(e) an antibody that binds to an expression product of NA group 3 nucleic acids,

(f) an antibody that binds to an expression product of NA group 5 nucleic acids,  
20

(g) an agent that binds to a complex of an HLA molecule and a fragment of an expression product of a NA group 1 nucleic acid,

(h) an agent that binds to a complex of an HLA molecule and a fragment of an expression product of a NA group 3 nucleic acid, and

(i) an agent that binds to a complex of an HLA molecule and a fragment of an  
25 expression product of a NA group 5 nucleic acid.

3. The method of claim 1, wherein the disorder is characterized by expression of a plurality of human cancer associated antigen precursors and wherein the agent is a plurality of  
30 agents, each of which is specific for a different human cancer associated antigen precursor, and wherein said plurality of agents is at least 2, at least 3, at least 4, at least 4, at least 6, at least 7, or at least 8, at least 9 or at least 10 such agents.

4. The method of claims 1-3, wherein the disorder is selected from the group consisting of small cell lung cancer, non-small cell lung cancer, melanoma, colon cancer, breast cancer, head and neck cancer, transitional cancer, leiomyosarcoma and synovial sarcoma.
5. The method of claim 1, wherein the nucleic acid molecule is selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12).
6. The method of claim 1, wherein the biological sample is isolated from a tissue selected from the group consisting of a non-brain, non-testis, non-prostate, non-small intestine and non-colon tissue.
7. A method for determining regression, progression or onset of a condition characterized by expression of abnormal levels of a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule, comprising  
monitoring a sample, from a patient who has or is suspected of having the condition, for a parameter selected from the group consisting of
  - (i) the protein,
  - (ii) a peptide derived from the protein,
  - (iii) an antibody which selectively binds the protein or peptide, and
  - (iv) cytolytic T cells specific for a complex of the peptide derived from the protein and an MHC molecule,as a determination of regression, progression or onset of said condition.
8. The method of claim 7, wherein the sample is a body fluid, a body effusion or a tissue.
9. The method of claim 7, wherein the step of monitoring comprises contacting the sample with a detectable agent selected from the group consisting of
  - (a) an antibody which selectively binds the protein of (i), or the peptide of (ii),
  - (b) a protein or peptide which binds the antibody of (iii), and
  - (c) a cell which presents the complex of the peptide and MHC molecule of (iv).
10. The method of claim 9, wherein the antibody, the protein, the peptide or the cell is

labeled with a radioactive label or an enzyme.

11. The method of claim 7, comprising assaying the sample for the peptide.

5 12. The method of claim 7, wherein the nucleic acid molecule is a NA Group 3 molecule or a NA Group 5 molecule.

13. The method of claim 7, wherein the nucleic acid molecule is selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3  
10 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12).

14. The method of claim 7, wherein the protein is a plurality of proteins, the parameter is a plurality of parameters, each of the plurality of parameters being specific for a different of the plurality of proteins, at least one of which is a cancer associated protein encoded by a NA  
15 Group 1 molecule.

15. The method of claim 7, wherein the protein is a plurality of proteins, at least one of which is encoded by SOX2 (SEQ ID NO:3) or ZIC2 (SEQ ID NO:5), and wherein the parameter is a plurality of parameters, each of the plurality of parameters being specific for a  
20 different of the plurality of proteins.

16. A pharmaceutical preparation for a human subject comprising  
an agent which when administered to the subject enriches selectively the presence of complexes of an HLA molecule and a human cancer associated antigen, and  
25 a pharmaceutically acceptable carrier, wherein the human cancer associated antigen is a fragment of a human cancer associated antigen precursor encoded by a nucleic acid molecule which comprises a NA Group 1 molecule.

17. The pharmaceutical preparation of claim 16, wherein the agent comprises a plurality of  
30 agents, each of which enriches selectively in the subject complexes of an HLA molecule and a different human cancer associated antigen, wherein at least one of the human cancer associated antigens is encoded by a NA Group 1 molecule.

18. The pharmaceutical preparation of claim 17, wherein the plurality is at least two, at least three, at least four or at least 5 different such agents.

19. The pharmaceutical preparation of claim 16, wherein the nucleic acid molecule is a  
5 NA Group 3 nucleic acid molecule.

20. The pharmaceutical preparation of claim 16, wherein the agent comprises a plurality of agents, at least one of which is a nucleic acid selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and  
10 SOX21 (SEQ ID NO:12), or an expression product thereof, each of which enriches selectively in the subject complexes of an HLA molecule and a different human cancer associated antigen.

21. The pharmaceutical preparation of claim 14, wherein the agent is selected from the  
15 group consisting of

(1) an isolated polypeptide comprising the human cancer associated antigen, or a functional variant thereof,

(2) an isolated nucleic acid operably linked to a promoter for expressing the isolated polypeptide, or functional variant thereof,

20 (3) a host cell expressing the isolated polypeptide, or functional variant thereof, and

(4) isolated complexes of the polypeptide, or functional variant thereof, and an HLA molecule.

22. The pharmaceutical preparation of claims 16-21, further comprising an adjuvant.  
25

23. The pharmaceutical preparation of claim 16, wherein the agent is a cell expressing an isolated polypeptide comprising the human cancer associated antigen or a functional variant thereof, and wherein the cell is nonproliferative.

30 24. The pharmaceutical preparation of claim 16, wherein the agent is a cell expressing an isolated polypeptide comprising the human cancer associated antigen or a functional variant thereof, and wherein the cell expresses an HLA molecule that binds the polypeptide.

25. The pharmaceutical preparation of claim 23 or 24, wherein the isolated polypeptide comprises a polypeptide encoded by a nucleic acid molecule selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12).
- 5
26. The pharmaceutical preparation of claim 16, wherein the agent is at least two, at least three, at least four or at least five different polypeptides, each coding for a different human cancer associated antigen or functional variant thereof, wherein at least one of the human cancer associated antigens is encoded by a NA Group 1 molecule.
- 10
27. The pharmaceutical preparation of claim 26, wherein the at least one of the human cancer associated antigens is a polypeptide encoded by a nucleic acid molecule selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12), or a fragment thereof.
- 15
28. The pharmaceutical preparation of claim 16, wherein the agent is a PP Group 2 polypeptide.
29. The pharmaceutical preparation of claim 16, wherein the agent is a PP Group 3 polypeptide or a PP Group 4 polypeptide.
- 20
30. The pharmaceutical preparation of claim 24, wherein the cell expresses one or both of the polypeptide and HLA molecule recombinantly.
- 25
31. The pharmaceutical preparation of claim 24, wherein the cell is nonproliferative.
32. A composition comprising  
an isolated agent that binds selectively a PP Group 1 polypeptide.
- 30
33. The composition of matter of claim 32, wherein the agent binds selectively a PP Group 2 polypeptide.
34. The composition of matter of claim 32, wherein the agent binds selectively a PP

Group 3 polypeptide.

35. The composition of matter of claim 32, wherein the agent binds selectively a PP Group 4 polypeptide.

5

36. The composition of matter of claim 32, wherein the agent binds selectively a PP Group 5 polypeptide.

10

37. The composition of claims 32-36, wherein the agent is a plurality of different agents that bind selectively at least two, at least three, at least four, or at least five different such polypeptides.

15

38. The composition of claim 37, wherein the at least one of the polypeptides is a polypeptide encoded by a nucleic acid molecule selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12), or a fragment thereof.

20

39. The composition of claims 32-36, wherein the agent is an antibody.

40. The composition of claim 37 wherein the agent is an antibody.

41. A composition of matter comprising  
a conjugate of the agent of claims 32-36 and a therapeutic or diagnostic agent.

25

42. A composition of matter comprising  
a conjugate of the agent of claim 37 and a therapeutic or diagnostic agent.

43. The composition of matter of claim 41, wherein the conjugate is of the agent and a therapeutic or diagnostic that is a toxin.

30

44. A pharmaceutical composition comprising an isolated nucleic acid molecule selected from the group consisting of NA Group 1 molecules and NA Group 2 molecules, and a pharmaceutically acceptable carrier.

45. The pharmaceutical composition of claim 44, wherein the isolated nucleic acid molecule comprises a NA Group 3 or NA Group 4 molecule.

5 46. The pharmaceutical composition of claim 44, wherein the isolated nucleic acid molecule comprises at least two isolated nucleic acid molecules coding for two different polypeptides, each polypeptide comprising a different human cancer associated antigen.

47. The pharmaceutical composition of claim 46, wherein at least one of the nucleic acid  
10 molecules is selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12).

48. The pharmaceutical composition of claims 44-47 further comprising an expression  
15 vector with a promoter operably linked to the isolated nucleic acid molecule.

49. The pharmaceutical composition of claims 44-47 further comprising a host cell  
recombinantly expressing the isolated nucleic acid molecule.

50. A pharmaceutical composition comprising  
20 an isolated polypeptide comprising a PP Group 1 or a PP Group 2 polypeptide, and a pharmaceutically acceptable carrier.

51. The pharmaceutical composition of claim 50, wherein the isolated polypeptide  
25 comprises a PP Group 3 or a PP Group 4 polypeptide.

52. The pharmaceutical composition of claim 50, wherein the isolated polypeptide  
comprises at least two different polypeptides, each comprising a different human cancer  
associated antigen.

30 53. The pharmaceutical composition of claim 52, wherein at least one of the polypeptides is a polypeptide encoded by a nucleic acid molecule selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12).

54. The pharmaceutical composition of claims 50-53, further comprising an adjuvant.
55. An isolated nucleic acid molecule comprising a NA Group 3 molecule.
56. An isolated nucleic acid molecule comprising a NA Group 4 molecule.
57. An isolated nucleic acid molecule selected from the group consisting of
- (a) a fragment of a nucleic acid molecule having a nucleotide sequence selected from the group consisting of nucleotide sequences set forth as SEQ ID NOs. 3-17, of sufficient length to represent a sequence unique within the human genome, and identifying a nucleic acid encoding a human cancer associated antigen precursor,
- (b) complements of (a),
- provided that the fragment includes a sequence of contiguous nucleotides which is not identical to any sequence selected from the sequence group consisting of
- (1) sequences having the GenBank accession numbers of Table 4,
- (2) complements of (1), and
- (3) fragments of (1) and (2).
58. The isolated nucleic acid molecule of claim 50, wherein the sequence of contiguous nucleotides is selected from the group consisting of:
- (1) at least two contiguous nucleotides nonidentical to the sequence group,
- (2) at least three contiguous nucleotides nonidentical to the sequence group,
- (3) at least four contiguous nucleotides nonidentical to the sequence group,
- (4) at least five contiguous nucleotides nonidentical to the sequence group,
- (5) at least six contiguous nucleotides nonidentical to the sequence group,
- (6) at least seven contiguous nucleotides nonidentical to the sequence group.
59. The isolated nucleic acid molecule of claim 57, wherein the fragment has a size selected from the group consisting of at least: 8 nucleotides, 10 nucleotides, 12 nucleotides, 14 nucleotides, 16 nucleotides, 18 nucleotides, 20, nucleotides, 22 nucleotides, 24 nucleotides, 26 nucleotides, 28 nucleotides, 30 nucleotides, 50 nucleotides, 75 nucleotides, 100 nucleotides, and 200 nucleotides.

60. The isolated nucleic acid molecule of claim 57, wherein the molecule encodes a polypeptide which, or a fragment of which, binds a human HLA receptor or a human antibody.

5

61. An expression vector comprising an isolated nucleic acid molecule of any of claims 55-60 operably linked to a promoter.

10

62. An expression vector comprising a nucleic acid operably linked to a promoter, wherein the nucleic acid is a NA Group 2 molecule.

63. An expression vector comprising a NA Group 1 or Group 2 molecule and a nucleic acid encoding an HLA molecule.

15

64. A host cell transformed or transfected with an expression vector of claim 61.

65. A host cell transformed or transfected with an expression vector of claims 62 or 63.

20

66. A host cell transformed or transfected with an expression vector of claim 61 and further comprising a nucleic acid encoding HLA.

67. A host cell transformed or transfected with an expression vector of claim 62 and further comprising a nucleic acid encoding HLA.

25

68. An isolated polypeptide encoded by the isolated nucleic acid molecule of claim 55 or claim 56.

69. A fragment of the polypeptide of claim 68 which is immunogenic.

30

70. An isolated polypeptide comprising a fragment of a polypeptide selected from the group consisting of ZIC2, SOX1, SOX2, SOX3 and SOX21 which is immunogenic, wherein the isolated polypeptide is not a full-length ZIC2, SOX1, SOX2, SOX3 or SOX21 polypeptide.

71. The polypeptide of claims 69 or 70, wherein the fragment, or a portion of the fragment, binds a HLA molecule or a human antibody.
- 5 72. An isolated fragment of a human cancer associated antigen precursor which, or a portion of which, binds a HLA molecule or a human antibody, wherein the precursor is encoded by a nucleic acid molecule that is a NA Group 1 molecule.
73. The fragment of claim 72, wherein the fragment is part of a complex with the HLA  
10 molecule.
74. The fragment of claim 73, wherein the fragment is between 8 and 12 amino acids in length.
- 15 75. An isolated polypeptide comprising a fragment of the polypeptide of claim 68 of sufficient length to represent a sequence unique within the human genome and identifying a polypeptide that is a human cancer associated antigen precursor.
76. A kit for detecting the presence of the expression of a human cancer associated antigen  
20 precursor comprising  
a pair of isolated nucleic acid molecules each of which consists essentially of a molecule selected from the group consisting of (a) a 12-32 nucleotide contiguous segment of the nucleotide sequence of any of the NA Group 1 molecules and (b) complements of ("a"), wherein the contiguous segments are nonoverlapping.
- 25 77. The kit of claim 76, wherein the pair of isolated nucleic acid molecules is constructed and arranged to selectively amplify an isolated nucleic acid molecule that is a NA Group 3 molecule.
- 30 78. A method for treating a subject with a disorder characterized by expression of a human cancer associated antigen precursor, comprising  
administering to the subject an amount of an agent, which enriches selectively in the subject the presence of complexes of a HLA molecule and a human cancer associated antigen,

effective to ameliorate the disorder, wherein the human cancer associated antigen is a fragment of a human cancer associated antigen precursor encoded by a nucleic acid molecule selected from the group consisting of

- (a) a nucleic acid molecule comprising NA group 1 nucleic acid molecules,
- 5 (b) a nucleic acid molecule comprising NA group 3 nucleic acid molecules, and
- (c) a nucleic acid molecule comprising NA group 5 nucleic acid molecules.

79. The method of claim 78, wherein the disorder is characterized by expression of a plurality of human cancer associated antigen precursors and wherein the agent is a plurality of  
10 agents, each of which enriches selectively in the subject the presence of complexes of an HLA molecule and a different human cancer associated antigen, wherein at least one of the human cancer associated antigens is encoded by a NA Group 1 molecule.

80. The method of claim 79, wherein at least one of the human cancer associated antigens  
15 is a polypeptide encoded by a nucleic acid molecule selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12), or a fragment thereof.

81. The method of claim 79, wherein the plurality is at least 2, at least 3, at least 4, or at  
20 least 5 such agents.

82. The method of claims 78-81, wherein the agent is an isolated polypeptide selected from the group consisting of PP Group 1, PP Group 2, PP Group 3, PP Group 4, and PP  
25 Group 5.

83. The method of claims 78-81, wherein the disorder is cancer.

84. The method of claims 82, wherein the disorder is cancer.

30 85. A method for treating a subject having a condition characterized by expression of a human cancer associated antigen precursor in cells of the subject, comprising:

- (i) removing an immunoreactive cell containing sample from the subject,
- (ii) contacting the immunoreactive cell containing sample to the host cell under

conditions favoring production of cytolytic T cells against a human cancer associated antigen which is a fragment of the precursor,

- (iii) introducing the cytolytic T cells to the subject in an amount effective to lyse cells which express the human cancer associated antigen, wherein the host cell is transformed or transfected with an expression vector comprising an isolated nucleic acid molecule operably linked to a promoter, the isolated nucleic acid molecule being selected from the group of nucleic acid molecules consisting of NA Group 1, NA Group 2, NA Group 3, NA Group 4, and NA Group 5.

86. The method of claim 85, wherein the host cell recombinantly expresses an HLA molecule which binds the human cancer associated antigen.

87. The method of claim 85, wherein the host cell endogenously expresses an HLA molecule which binds the human cancer associated antigen.

88. A method for treating a subject having a condition characterized by expression of a human cancer associated antigen precursor in cells of the subject, comprising:

- (i) identifying a nucleic acid molecule expressed by the cells associated with said condition, wherein said nucleic acid molecule is a NA Group 1 molecule;
- (ii) transfecting a host cell with a nucleic acid selected from the group consisting of (a) the nucleic acid molecule identified, (b) a fragment of the nucleic acid identified which includes a segment coding for a human cancer associated antigen, (c) deletions, substitutions or additions to (a) or (b), and (d) degenerates of (a), (b), or (c);

(iii) culturing said transfected host cells to express the transfected nucleic acid molecule, and;

(iv) introducing an amount of said host cells or an extract thereof to the subject effective to increase an immune response against the cells of the subject associated with the condition.

89. The method of claim 88, wherein the nucleic acid molecule is selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12).

90. The method of claim 88, further comprising identifying an MHC molecule which presents a portion of an expression product of the nucleic acid molecule, wherein the host cell expresses the same MHC molecule as identified and wherein the host cell presents an MHC binding portion of the expression product of the nucleic acid molecule.

5

91. The method of claim 88, wherein the immune response comprises a B-cell response or a T cell response.

10

92. The method of claim 91, wherein the response is a T-cell response which comprises generation of cytolytic T-cells specific for the host cells presenting the portion of the expression product of the nucleic acid molecule or cells of the subject expressing the human cancer associated antigen.

15

93. The method of claim 88, wherein the nucleic acid molecule is a NA Group 3 molecule.

94. The method of claims 88 or 90, further comprising treating the host cells to render them non-proliferative.

20

95. A method for treating or diagnosing or monitoring a subject having a condition characterized by expression of an abnormal amount of a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule, comprising

administering to the subject an antibody which specifically binds to the protein or a peptide derived therefrom, the antibody being coupled to a therapeutically useful agent, in an amount effective to treat the condition.

25

96. The method of claim 95, wherein the antibody is a monoclonal antibody.

97. The method of claim 96, wherein the monoclonal antibody is a chimeric antibody or a humanized antibody.

30

98. A method for treating a condition characterized by expression in a subject of abnormal amounts of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule, comprising

administering to a subject a pharmaceutical composition of any one of claims 16-31 and 44-54 in an amount effective to prevent, delay the onset of, or inhibit the condition in the subject.

5 99. The method of claim 98, wherein the condition is cancer.

100. The method of claim 98, further comprising first identifying that the subject expresses in a tissue abnormal amounts of the protein.

10 101. The method of claim 99, further comprising first identifying that the subject expresses in a tissue abnormal amounts of the protein.

102. A method for treating a subject having a condition characterized by expression of abnormal amounts of a protein encoded by a nucleic acid molecule that is a NA Group 1  
15 nucleic acid molecule, comprising  
(i) identifying cells from the subject which express abnormal amounts of the protein;  
(ii) isolating a sample of the cells;  
(iii) cultivating the cells, and  
(iv) introducing the cells to the subject in an amount effective to provoke an immune  
20 response against the cells.

103. The method of claim 102, further comprising rendering the cells non-proliferative, prior to introducing them to the subject.

25 104. A method for treating a pathological cell condition characterized by aberrant expression of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule, comprising  
administering to a subject in need thereof an effective amount of an agent which inhibits the expression or activity of the protein.

30

105. The method of claim 104, wherein the agent is an inhibiting antibody which selectively binds to the protein and wherein the antibody is a monoclonal antibody, a chimeric antibody, a humanized antibody or an antibody fragment.

106. The method of claim 104, wherein the agent is an antisense nucleic acid molecule which selectively binds to the nucleic acid molecule which encodes the protein.

5 107. The method of claim 104, wherein the nucleic acid molecule is a NA Group 3 nucleic acid molecule.

108. The method of claim 104, wherein the nucleic acid molecule is selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5),  
10 SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12).

109. A composition of matter useful in stimulating an immune response to a plurality of a proteins encoded by nucleic acid molecules that are NA Group 1 molecules, comprising  
a plurality of peptides derived from the amino acid sequences of the proteins, wherein  
15 the peptides bind to one or more MHC molecules presented on the surface of the cells which express an abnormal amount of the protein.

110. The composition of matter of claim 109, wherein at least a portion of the plurality of peptides bind to MHC molecules and elicit a cytolytic response thereto.

20 111. The composition of matter of claim 109, wherein at least one of the proteins is encoded by a nucleic acid molecule selected from the group consisting of SOX2 (SEQ ID NO:3), SOX1 (SEQ ID NO:4), ZIC2 (SEQ ID NO:5), SOX3 (SEQ ID NO:11) and SOX21 (SEQ ID NO:12).

25 112. The composition of matter of claim 110, further comprising an adjuvant.

113. The composition of matter of claim 112, wherein said adjuvant is a saponin, GM-CSF, or an interleukin.

30 114. The composition of matter of claim 109, further comprising at least one peptide useful in stimulating an immune response to at least one protein which is not encoded by nucleic acid molecules that are NA Group 1 molecules, wherein the at least one peptide binds to one

or more MHC molecules.

115. An isolated antibody which selectively binds to a complex of:

- 5 (i) a peptide derived from a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule and
- (ii) and an MHC molecule to which binds the peptide to form the complex, wherein the isolated antibody does not bind to (i) or (ii) alone.

116. The antibody of claim 115, wherein the antibody is a monoclonal antibody, a chimeric  
10 antibody, a humanized antibody, or a fragment thereof.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
22

**Abstract**

Cancer associated antigens have been identified by autologous antibody screening of libraries of nucleic acids expressed in small cell lung cancer cells using antisera from cancer patients. The invention relates to nucleic acids and encoded polypeptides which are cancer associated antigens expressed in patients afflicted with small cell lung cancer. The invention provides, *inter alia*, isolated nucleic acid molecules, expression vectors containing those molecules and host cells transfected with those molecules. The invention also provides isolated proteins and peptides, antibodies to those proteins and peptides and cytotoxic T lymphocytes which recognize the proteins and peptides. Fragments of the foregoing including functional fragments and variants also are provided. Kits containing the foregoing molecules additionally are provided. The molecules provided by the invention can be used in the diagnosis, monitoring, research, or treatment of conditions characterized by the expression of one or more cancer associated antigens.

15

[illegible]

276

Variable	Mean	SD	Min	Max
Age	38.5	12.5	18	65
Gender	0.5	0.5	0	1
Marital status	0.7	0.5	0	1
Education	12.5	2.5	9	16
Income	3500	1500	1000	8000
Health status	0.8	0.4	0	1
Exercise frequency	2.5	1.5	0	5
Stress level	4.5	1.5	1	7
Sleep quality	3.5	1.5	1	7
Work satisfaction	4.0	1.5	1	7
Life satisfaction	5.5	1.5	1	7
Depression score	1.5	1.5	0	5
Anxiety score	1.5	1.5	0	5
Resilience score	4.0	1.5	1	7
Self-efficacy score	4.5	1.5	1	7
Optimism score	5.0	1.5	1	7
Gratitude score	5.5	1.5	1	7
Forgiveness score	5.0	1.5	1	7
Empathy score	5.5	1.5	1	7
Compassion score	5.0	1.5	1	7
Kindness score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Shamelessness score	5.0	1.5	1	7
Modesty score	5.5	1.5	1	7
Generosity score	5.0	1.5	1	7
Patience score	5.5	1.5	1	7
Humility score	5.0	1.5	1	

	<110> Gure, Ali	
	Stockert, Elisabeth	
5	Scanlan, Matthew J.	
	Jager, Dirk	
	Old, Lloyd J.	
	Chen, Yao-Tseng	
10	<120> SMALL CELL LUNG CANCER ASSOCIATED	
	ANTIGENS AND USES THEREOF	
	<130> L0461/7073	
15	<160> 22	
	<170> FastSEQ for Windows Version 3.0	
	<210> 1	
20	<211> 29	
	<212> DNA	
	<213> Homo sapiens	
	<400> 1	
25	catgaatatg aacatgggta tgaacatgg	29
	<210> 2	
	<211> 23	
	<212> DNA	
30	<213> Homo sapiens	
	<400> 2	
	tcgcagccct caaactcaca ctg	23
35	<210> 3	
	<211> 1085	
	<212> DNA	
	<213> Homo sapiens	
40	<400> 3	
	cacagcgccc gcatgtacaa catgatggag acggagctga agccgcgggg cccgcagcaa	60
	acttcggggg gcgggggcgg caactccacc gcggcgggcg ccggcgggcaa ccagaaaaac	120
	agcccggaacc gcgtcaagcg gcccatgaat gccttcatgg tgttgtcccg cgggcagcgg	180
	cgcaagatgg cccaggagaa cccaagatg cacaactcgg agatcagcaa gcgcctgggc	240
45	gccgagtgga aactttttgtc ggagacggag aagcgggcgt tcatcgacga ggctaagcgg	300
	ctgcgagcgc tgcacatgaa ggagcaccgc gattataaat accggccccg gcggaaaacc	360
	aagacgctca tgaagaagga taagtacacg ctgcccggcg ggctgctggc ccccgggcggc	420
	aatagcatgg cgagcggggg cgggggtggc gccggcctgg gcgcggggcg gaaccagcgc	480
	atggacagtt acgcgcacat gaacggctgg agcaacggca gctacagcat gatgcaggac	540
50	cagctgggct accgcgacga cccgggectc aatgcgcacg gcgcagcgca gatgcagccc	600
	atgcaccgct acgacgtgag cgccctgcag tacaactcca tgaccagctc gcagacctac	660
	atgaacggct cgcccaccta cagcatgtcc tactcgagc agggcacccc tggcatggct	720
	cttggtctca tgggttcggg ggtcaagtcc gaggccagct ccagcccccc tgttggttacc	780
	tcttctctcc actccagggc gccctgccag gccggggacc tccgggacat gatcagcatg	840
55	tatctccccg gcgcgaggt gccggaacc gcgcggggcg gcagacttca catgtcccag	900
	cactaccaga cgggcccggg gcccggcacg gccattaacg gcacactgcc cctctcacac	960
	atgtgagggc cggacagcga actggagggg ggagaaattt tcaaagaaaa acgagggaaa	1020
	tgggaggggt gcaaaagagg agagtaagaa acagcatgga gaaaaccogg tacgtctaaa	1080
	aaaaa	1085
60	<210> 4	
	<211> 4091	
	<212> DNA	

<213> Homo sapiens

<400> 4

	ccggccgtct	atgctccagg	ccctctctc	gcgggtgccg	tgaacccgcc	agccgccccg	60
5	atgtacagca	tgatgatgga	gaccgacctg	cactcgcccc	gcggcgccca	ggccccacg	120
	aacctctcgg	gccccgcgg	ggcgggcggc	ggcgggggcg	gaggcggggg	cgcgggcggc	180
	ggcgggggcg	ccaaggccaa	ccaggacctg	gtcaaacggc	ccatgaacgc	cttcatggtg	240
	tgggtcccgcg	ggcagcggcg	caagatggcc	caggagaacc	ccaagatgca	caactcggag	300
	atcagcaagc	gcctgggggc	cgagtggaa	gtcatgtccg	aggccgagaa	gcggccgttc	360
10	atcgacgagg	ccaagcggct	gcgcgcgctg	cacatgaagg	agcacccgga	ttacaagtac	420
	cggccgcgcc	gcaagaccaa	gacgctgctc	aagaaggaca	agtactcgct	ggccggcggg	480
	ctcctggcgg	ccggcgcggg	tggcgcggc	gcggctgtgg	ccatgggcgt	gggcgtgggc	540
	tggggcgcg	cgcccgctgg	ccagcgctg	gagagcccag	gcggcgcggc	ggcgcgcgcg	600
	tacgcgcacg	tcaacggctg	ggccaacggc	gcctaccccg	gctcgggtgg	ggccgcggcg	660
15	gcgcgcgcgg	ccatgatgca	ggaggcgag	ctggcctacg	ggcagcacc	cgcgcggggc	720
	ggcgcgccac	cgcaccgcac	ccggcgccac	ccgcacccgc	accaccgca	cgcgccaccg	780
	cacaacccgc	agcccatgca	ccgctacgac	atggcgcgcg	tgcagtacag	ccccatctcc	840
	aactcgcagg	gctacatgag	cgcgtcgccc	tcgggctacg	gcggcctccc	ctacggcgcc	900
	gcggccgcgc	ccgcgcgcgc	gcaccagaac	tcggccgtgg	cgccggcggc	ggcgcgcgcg	960
20	gcgcgctcgt	cgggcgccct	ggcgcgctg	ggctctctgg	tgaagtcgga	ggccagcggc	1020
	agcccccgct	ccccagcgca	ctcgcgggcg	ccgtgccccg	gggacctgcg	cgagatgac	1080
	agcatgtact	tgcgcgcgg	cgagggggcg	gaccgcggcg	cggcagcagc	ggcgcgcgcg	1140
	cagagccggc	tgcactcgct	gcgcagcac	taccagggcg	cgggcgcggg	cgtgaacggc	1200
	acggtgcccc	tgacgcacat	ctagcgctt	cgggacgccc	gggactctgc	ggcgcggaac	1260
25	cacgagctcg	cgcccgcgcg	ccggctcccc	ccccgccccg	gcgcggcggtg	gcttttgtat	1320
	cagacgttcc	cacattcttg	tcaaaaggaa	aatactggag	acgaacgccc	ggtgacgcgt	1380
	gtccccact	caccttcccc	ggagaccctg	gcgaccgccc	ggcgctgaca	ccagacttgg	1440
	tttagactga	acttcgggtg	tttcttgaga	cttttgtaca	gtattttatca	cctacggagg	1500
	aagcgggaagc	gttttctttg	ctcgagggga	caaaaaagtc	aaaacgaggc	gagaggcgaa	1560
30	gcccactttt	gtataccggc	cggcgcgctc	actttctctc	gcgttgcttc	cggaacggcg	1620
	gcaccgcggc	agccccagtg	acgcggagct	cgtcgcatct	gttataaatg	tagtaaggca	1680
	ggccaagca	cttacaagtt	ttttgtagtt	gttaccgctc	ttttgggttg	gtttgttaat	1740
	ttatacaaa	agattaccac	caccaccccc	tccttcagac	ggcgaggtta	tattctgggt	1800
	tttgtaaaac	tttatgtatc	tgagcatttc	catttttttt	tttggttttt	gtattatttc	1860
35	ttgtaaatgc	atttgtaaaa	attttatttt	cggcgttgca	atgcggggag	gagaagtcag	1920
	attatgtaca	tagttttcta	aaaagccttt	cttctaaaaa	cgaaaaaaga	ccccaccca	1980
	aatgttttcg	agtcaacaaa	tttaagagac	agagcccat	ttctccataa	atttgtaaca	2040
	tgcctatttt	tatgtgcatg	ttttatgagt	tcaaaatgca	atgagggaaa	tctgacaggg	2100
	aaattatctg	tatgaactaa	aagtaaggga	accgggggaa	tgggaggaca	ggatttttca	2160
40	aggaaccttt	ttcaatgaaa	gagaaggga	ttaaaacct	taggttat	tgtagagctg	2220
	agtgttaata	cgggcgagaa	aataaaagta	tcttctgctc	cggtctgttc	actgcggacg	2280
	gctggggctg	ctgcgcgtta	ccttgctgca	acngggcgcc	ttccacctgg	ctgggggtct	2340
	gcgccacagt	ttggtccaga	ngwgggagga	ggaagggaag	acccagtggt	tgggaccttg	2400
	gaccaggcca	tggatgaagg	acaaagacca	gggcaggtca	cggttttccc	aattccccag	2460
45	caattaagat	ttcgagcaga	atztatctaa	atgtgtttca	aggaaacaca	atcgctgaac	2520
	caaaacgtac	tgcagccgan	ccccctccgt	ccatcctctg	cccccccccc	tggcttcttt	2580
	ctcttgggaa	aacggggcaa	ataattgtgc	tggattctca	cacacacaga	aatactgacc	2640
	atcacccctc	cccgctgaa	ctgggatgca	agttgctaac	cgatgtgaac	gcaaaatgcc	2700
	ttgttcatta	ttcctgacga	gatcttgagg	ttgtttgatg	ctttaaattt	tttaattata	2760
50	ttattttcta	ggtgtttatt	ggtacattgc	agtttttttt	ttgaaattta	aaaattttctg	2820
	taaaactttg	tcttcaagta	atctgacagc	attaaatatt	gcatttaaaa	attatactgt	2880
	agcaaataca	tttaaaaatt	aatcacacgc	ttaagatgaa	attataattt	tggaaaaaaa	2940
	aaacacttga	agccagatg	gaaatacgtt	tatttcagca	gccttaggtt	ttccctcgct	3000
	ttctcaacac	ccttctctgt	cctggagtat	ggactgtccg	tcctaaaagt	agcctatgct	3060
55	ataagtttaa	tgagaaccca	atcagccctg	cattcgagaa	tagcttttaag	tataatgctg	3120
	atctgacaat	tgacgtgtaa	tttgggaagt	cattttgata	attttgctta	aaccactcat	3180
	tcgttaaagt	gattacaaaa	aagttcaaga	atgatgtcca	ctgctttcta	acaagataat	3240
	aaaccccccc	cctcttttct	ttttctttat	ttttatttct	tttagctatt	tgatcctttc	3300
	tgaagcagtt	gtttctggaa	gagtctgtgc	gcccattggat	ggctgagcac	cactacgact	3360
60	tagtccggga	taagggcctc	cccagtcctc	tccgggagat	gatttgggaa	attttataat	3420
	gcttggtctg	ttaactcacc	gggaccttga	gggtccaatg	ggacctgag	ggttttctct	3480
	gaaatatata	aacttaaagg	actctctctg	aggttctttg	actgacgtcc	actctcagtc	3540
	tggccctctg	gctccctctg	gtgtaccctg	gagtttctgt	gtccaattgt	tggcatctag	3600

	gtcttggctc	aagattagga	tgtgggcccc	acttttagagg	cacagactat	gaaaagctga	3660
	gttagtgccg	ccgggacgcc	aggcaagcag	cttttacagt	ttggcatctt	attgcagggtg	3720
	cttcgtgcac	agtcagctga	aatagccaat	gccagggtgct	ccaaccacct	tatttcottg	3780
	ttttgttgat	tagaacaaca	cagaaaaaag	caaataataa	tttttaatga	ctccatttaa	3840
5	aaatatcaca	gggtgggggc	aaggaaatta	gctgagattc	atctcaggat	tgagattcta	3900
	tcccccttc	cccgcccca	gcagtgtcgc	tccaattcaa	attagtggag	aaaagattac	3960
	agtaggccct	gagccgactg	tgaattcggg	gcttggccaa	ggtaacactc	atcgtattca	4020
	cggagraaat	actatatgat	gatagttatt	atattatatg	acgacttcat	tcacttccca	4080
	aatcacaggg	t					4091

10

<210> 5  
<211> 1602  
<212> DNA  
<213> Homo sapiens

15	<400> 5						
	atgctcctgg	acgcgggtcc	gcagttcccg	gccatcgggg	tgggcagctt	cgcgcgccac	60
	catcaccact	ccgcgcgggc	ggcggcggcg	gctgccgcgc	agatgcagga	ccgtgaactg	120
	agcctggcgg	cgggcgagaa	cggtctcggt	gattccgcgc	ccgcgcacat	gggagccttc	180
20	aagctcaacc	cgggcgcgca	cgagctgtcc	ccgggccaga	gctcggcggt	cacgtcgcag	240
	ggccccggcg	cctaccccg	ctccgctgcg	gctgccgctg	cgcccgccagc	gctcggggccc	300
	cacgcgcgcg	acgttggctc	ctactctggg	ccgcccttca	actccacccg	ggacttcctg	360
	ttccgcagcg	cgcggtcttc	ggggacttcg	gcgccggggcg	gcgggcagca	cgggctgttc	420
	gggcggggcg	cgggcggcct	gcaccacgcg	cactcggacg	cgcagggcca	cctcctcttc	480
25	ccgggcctgc	cagagcagca	cgggccgcac	ggctcgcaga	atgtgctcaa	cgggcagatg	540
	cgcctcgggc	tgcccggcga	ggtgttcggg	cgctcggagc	aataccgcca	ggtggccagc	600
	ccgcggaccg	acccctactc	ggcggcgcaa	ctccacaacc	agtacggccc	catgaatatg	660
	aacatgggta	tgaacatggc	agcagccgcg	gcccaccacc	accaccacca	ccaccaccac	720
	cccggtgcct	ttttccgcta	tatgcggcag	cagtgcacat	agcaggagct	aatctgcaag	780
30	tggatcgacc	ccgagcaact	gagcaatccc	aagaagagct	gcaacaaaac	tttcagcacc	840
	atgcacgagc	tggtgacaca	cgtctcgggt	gagcacgtcg	gcggcccgga	gcagagcaac	900
	cacgtctgct	tctgggagga	gtgtccgcgc	gagggcaagc	ccttcaaggc	caaatacaaa	960
	ctggtcaacc	acatccgcgt	gcacacaggc	gagaaacct	tcccctgccc	cttcccgggc	1020
	tgtggcaaa	tcttcgcgcg	ctccgagaac	ctcaagatcc	acaaaaggac	ccacacaggg	1080
35	gagaagccgt	tccagtgtga	gtttgagggc	tgcgaccggc	gcttcgccaa	cagcagcgac	1140
	aggaagaagc	acatgcacgt	ccacacctcc	gataagccct	atctctgcaa	gatgtgcgac	1200
	aagtccctaca	cgcaccccag	ctcgtcgcgg	aagcacatga	aggtccatga	gtcctccccg	1260
	cagggttctg	aatcctcccc	ggcgccagc	tccggctatg	agtcgtccac	gcccccgggg	1320
	ctggtgtccc	ccagcgccga	gccccagagc	agctccaacc	tgtcccagc	ggcggcggca	1380
40	gcggcgccgg	cggctgcggc	ggcggcggcc	gcggtgtccg	cgggtgcaccg	gggcggaggc	1440
	tcgggcagtg	gcggcgcggg	aggcggtcca	ggcggcggca	gcggcagtg	cgggggcggc	1500
	ggcgggcgcg	gcggcggggg	cgcggcgagc	tctggcgggg	gcagcgggac	agccgggggt	1560
	cacagcggcc	tctcctccaa	cttcaatgaa	tggtacgtgt	ga		1602

45

<210> 6  
<211> 1322  
<212> DNA  
<213> Homo sapiens

50	<400> 6						
	ggaattccgg	gcgcggttgt	gagtagtacc	gggagtgggg	tgatcccggg	ctaggggagc	60
	gcggcgcccc	atcgggctta	gtcggagctc	cgaagggagt	gactaggaca	cccgggtggg	120
	ctacttttct	tccgggtgctt	ttgctttttt	tttccttttg	gctcgggctg	agtgtcgccc	180
	actgagcaaa	gattccctcg	taaaacccag	agcgaccctc	ccgtcaattg	ttgggctcgg	240
55	gagtgtcgcg	gtgccccgag	cgcgcggggc	gcggaggcaa	agggagcgga	gccggccgcg	300
	gacggggccc	ggagcttgcc	tgctccctc	gctcgcccca	gcgggttcgc	tcgcgtagag	360
	cgcagggcgc	gcgcgatgaa	ggcggtgagc	ccggtgcgcc	cctcgggccc	caaggcgccg	420
	tcgggctgcg	gcggcgggga	gctggcgctg	cgctgcctgg	ccgagcacgg	ccacagcctg	480
	ggtggctccg	cagccgcggc	ggcggcggcg	gcggcagcgc	gctgtaaggc	ggccgaggcg	540
60	gcggccgacg	agccgcgcgt	gtgcctgcag	tgcgatatga	acgactgcta	tagccgcctg	600
	cggaggctgg	tgcccacccat	cccgcocaa	aagaaagtca	gcaaagtgga	gatcctgcag	660
	cacgttatcg	actacatcct	ggacctgcag	ctggcgctgg	agacgcaccc	ggcctgctg	720
	aggcagccac	caccgcccgc	gccgccacac	caccgcggcg	ggacctgtcc	agccgcgcgc	780

	ccgcggaccc	cgctcactgc	gctcaacacc	gacccggccg	gcgcggtgaa	caagcagggc	840
	gacagcattc	tgtgccgctg	agccgcgctg	tccaggtgtg	cgcccgccctg	agcccgagcc	900
	aggagcacta	gagaggagg	gggaagagca	gaagttagag	aaaaaaagcc	accggaggaa	960
	aggaaaaaac	atcggccaac	ctagaaacgt	tttcattcgt	cattccaaga	gagagagagg	1020
5	aaagaaaaat	acaactttca	ttctttcttt	gcacgttcat	aaacattcta	catacgtatt	1080
	ctctttttgtc	tcttcattta	taactgctgt	gaattgtaca	ttttctgtgtt	ttttggagggt	1140
	gcagttaaac	ttttaagctt	aagtgtgaca	ggactgataa	atagaagatc	aagagtagat	1200
	ccgacttttag	aagcctactt	tgtgaccaag	gagctcaatt	tttgttttga	agctttacta	1260
	atctaccaga	gcattgtaga	tatttttttt	ttacatctat	tgttttaaat	agccggaatt	1320
10	cc						1322

<210> 7

<211> 2389

<212> DNA

15 <213> Homo sapiens

<400> 7

	cggtccagcg	ggggccgagg	ccatgttccc	ggtgtttcct	tgcacgctgc	tggccccccc	60
	cttccccgtg	ctgggacctg	actcccgggg	ggtgggcggc	ctcatgaact	ccttccccgc	120
20	acctcaggg	cacgcccaga	acccctgca	ggtcggggct	gagctccagt	cccgtttctt	180
	tgcctccag	ggctgcgccc	agagtccatt	ccaggcccg	ccggcgcccc	cgccccgcgc	240
	ccaggccccg	cgggccgagc	ccctccaggt	ggacttgctc	ccggtgctcg	ccgcgcgcca	300
	ggagtcgcgc	ggggtgctg	cgcccgctgc	cgccgctgct	gccgcgctcg	ctgcgcgcgc	360
	ccgggcccc	gccgcgcgct	ctacgggtgga	cacagcggcc	ctgaagcagc	ctccggcgcc	420
25	ccctccgcca	cccccgccag	tgtcggcgcc	cgccggccgag	gccgcgcccc	ccgcctccgc	480
	cgccactatc	gccgcggcgg	cgccaccgc	cgtcgtagcc	ccaacctcga	cggtcgcggt	540
	ggccccgggtc	ggtctgtcct	tggagaagaa	gacaaagagc	aagggggccct	acatctgcgc	600
	tctgtgcgc	aaggagttca	agaacggcta	caatctccgg	aggcacgaag	ccatccacac	660
	gggagccaa	gcccggccggg	tccctcggg	tgctatgaag	atgccgacca	tgggtgcccc	720
30	gagcctcctg	agcgtgcccc	agctgagcgg	agccggcggg	ggagggggag	aggcgggtgc	780
	cgccggcgcc	cctcccgag	tggccgcgg	tggcgtggtg	accacgaccg	cctcggggaa	840
	gcgcctccgg	aagaacctatg	cctgcgagat	gtgtggcaag	gccttccgcg	acgtctacca	900
	cctgaaccga	cacaagctgt	cgcactcgga	cgagaagccc	taccagtgcc	cggtgtgcca	960
	gcagcgcttc	aagcgcaagg	accgcatgag	ctaccacgtg	cgctcacatg	acggcgctgt	1020
35	gcacaagcc	tacaactgct	cccactgtgg	caagagcttc	tcccggcccg	atcacctcaa	1080
	cagtcacgtc	agacaagtgc	actcaacaga	acggcccttc	aaatgtgaga	aatgtgaggc	1140
	agctttcgcc	acgaaggatc	ggctgcgggc	gcacacagta	cgacacgagg	agaaagtgcc	1200
	atgtcacgtg	tgtggcaaga	tgtgagctc	ggcttatatt	tccgaccaca	tgaagggtgca	1260
	cagccaggg	cctcaccatg	tctgtgagct	ctgcaacaaa	ggtactggtg	aggtttgctc	1320
40	aatggcgggc	gcagcggcag	cgccggcagc	ggcagcagcg	gcagcagtag	cagccccctc	1380
	cacagctgtg	ggctccctct	cgggggcgga	gggggtgcct	gtgagctctc	agccacttcc	1440
	ctcccaaccc	tgggtgagctc	caagttgggt	gcgggggaga	ggggagaatg	gagtagagtc	1500
	ccttgggtaca	agctcctctc	ccccctcttt	tcccaccaac	tcctatttcc	ctaccaacca	1560
	aggagcctcc	agaaggaaa	gaggaagaaa	tgttttctta	ggggaattcg	ctagggtttta	1620
45	acgatttgc	tctcctgctc	ctcttctatc	agacctgacc	ccacacaaac	ctgtccccctc	1680
	ggttgtgttg	aagtccccctg	gacagtgggc	aggggtggca	gaggacacga	gcagccactg	1740
	cccgtaaccc	ctctcctctc	tgtaaagcca	tgccctgtct	tcccaggac	ttgtgagcct	1800
	cttccctcga	cggtcctctt	ctctccttcc	agtcctctcc	ccctgctgtc	tgcagcccc	1860
	ccccgggggag	ttggtgcttt	cttttctttt	tttttttttt	ttccaggggg	agggaggaga	1920
50	ggaaggagg	ggatcagagc	tgtcccaaag	agggaaagcg	gtgaggtttg	aggaggggca	1980
	gaagcagggc	cggcaaagg	tgtaccttca	taagtggtga	tccgggggtt	ggggtcaggc	2040
	cctgaacatc	gtcctacttg	agaatctgtc	aggggaaaaa	gtcaagggga	gcaggaggaa	2100
	gagccaggag	ggccagaggc	agagaagaga	tggagtctta	ggggccagg	tgagccagg	2160
	gtccaggggc	tagagggtgt	tctggggggg	ggggaatgca	gccagtgtcc	ccctccccctc	2220
55	ttccacccca	gctccagccc	tgggtcttgtc	ttttctctcc	tcttccccac	gacagaagaa	2280
	gttgtggccc	tggcatgtca	tgtgttccct	gtgtccctcg	catgtacccc	accctccacc	2340
	ccttcctttt	gcgcgggaacc	cattacaata	aatttttaaat	aaaatcctg		2389

<210> 8

<211> 1860

<212> DNA

60 <213> Homo sapiens

<400> 8

	gggacgtgag	cgctgcgcc	caccgggcta	gaccgggcgc	catcatgctg	cttctgccaa	60
	gcgccgcgga	cggccggggc	accgccatca	cccacgctct	gacctctgcc	tctacactct	120
	gtcaagttga	acctgtggga	agatggtttg	aagcttttgt	taagaggaga	aacagaaatg	180
5	cttctgcctc	ttttcaggaa	ctggaggata	agaaaagagt	atccgaggaa	tcagaagatg	240
	aagaattgca	gttggaagag	tttcccatgc	tgaaaacact	tgatcccaa	gactggaaga	300
	accaagatca	ttatgcagtt	cttggacttg	gccatgtgag	atacaaggct	acacagagac	360
	agatcaaagc	agctcataaa	gcaatggttt	taaaacatca	cccagacaaa	cggaaagcag	420
	ctggtgaacc	aataaaaagaa	ggagataaatg	actacttcac	ttgcataact	aaagcttatg	480
10	aaatgttatc	tgatccagtg	aaaagacgag	cattttaacag	tgtagatcct	acttttgata	540
	actcagttcc	ttctaaaagt	gaagcaaagg	ataattttctt	cgaagtgttt	accccagtg	600
	ttgaaaggaa	ttccagatgg	tcaaataaaa	aaaatgttcc	taaacttggg	gatattgaatt	660
	catcatttga	agatgtagat	atattttatt	ctttctggta	taattttgat	tcttggagag	720
	aattttctta	tttagatgaa	gaagaaaaag	aaaaagcaga	atgtcgtgat	gagaggagat	780
15	ggattgaaaa	gcagaacgga	gcaacaagag	cacaaagaaa	aaaagaagaa	atgaacagaa	840
	taagaacatt	agttgacaat	gcatacagct	gtgatccaag	gataaaaaag	ttcaagggaag	900
	aagaaaaagc	caagaaagaa	gcagaaaaga	aagcaaaagc	agaagctaaa	cgggaaggagc	960
	aagaagctaa	agaaaaacaa	agacaagctg	aattagaagc	tgctcgggta	gctaaggaga	1020
	aagaagagga	ggaagtcaga	cagcaagcat	tgctggcaaa	gaaggaaaaa	gatattccaga	1080
20	aaaaagccat	taagaaggaa	aggcaaaaac	ttcgaaactc	atgcaagata	gaagaaataa	1140
	atgagcaaat	cagaaaagag	aaagagggaag	ctgaggctcg	tatgcgacaa	gcattctaaga	1200
	acacagagaa	atcaactggg	ggagggtgaa	atggaaagtaa	aaattgggtca	gaagatgatac	1260
	tacaattact	aattaaagct	gtgaatctgt	tcctgtctag	aacaaattca	agatgggaag	1320
	ttattgctaa	ttacatgaac	atacattctt	cctctggagt	caaaaagaact	gccaaagatg	1380
25	ttattggcaa	agcaaagagt	ctccaaaaac	ttgacctca	tcaaaaagat	gacataaata	1440
	aaaaggcatt	tgataagttc	aaaaaagaac	atggagtggg	acctcaagca	gacaacgcaa	1500
	cgccttcaga	acgatttgaa	ggtccatata	cagacttcac	cccttgga	acagaagaac	1560
	agaagctttt	ggaacaagct	ttgaaaacat	accagtaaaa	tacacctgaa	agatgggaaa	1620
	aaatagcaga	agcgggtgct	ggcaggacaa	agaaggactg	catgaaacga	tacaagggaac	1680
30	ttgtcgagat	ggtaaaaagca	aagaaagctg	ctcaagaaca	agtgtgaat	gcaagtagag	1740
	ccaagaaatg	acaactcttg	ttgtgtgtgc	atttttataa	taaaactgaa	aatactgtaa	1800
	acattttcat	tcttaaaatt	atactcatgg	taataatttg	aaagtaaaaa	aaaaaaaaaa	1860

<210> 9

35	<211> 2291
	<212> DNA
	<213> Homo sapiens

<400> 9

40	gaattcctga	ctgccacagg	tgtacaggaa	acatttgtct	tttgttgctg	gaaagctgct	60
	caaatcaaa	aacatttact	gaagtcaaa	tggtgcgcc	ctacatctct	caatgtgggt	120
	cgaataatta	catcagagct	ctatcgatca	ctgggagatg	tcctccgtga	tggtgatgcc	180
	aaggcttttg	tgcgctctga	ctttcttctg	gtgtatggg	atgtcatctc	aaacatcaat	240
	atcaccagag	cccttgagga	acacaggttg	agacggaagc	tagaaaaaaa	tgtttctgtg	300
45	atgacgatga	tcttcaagga	gtcatcccc	agccacccaa	ctcgttgcca	cgaagacaat	360
	gtggtagtgg	ctgtggatag	taccacaaac	agggttctcc	attttcagaa	gaccagggtg	420
	ctccggcggt	ttgcatttcc	tctgagcctg	tttcagggca	gtagtgtatg	agtggagggt	480
	cgatatgatt	tactggattg	tcatatcagc	atctgttctc	ctcagggtggc	acaactcttt	540
	acagaaaact	ttgactacca	aactcgagat	gactttgtgc	gagggtctct	agtgaatgag	600
50	gagatcctag	ggaaccagat	ccacatgcac	gtaacagcta	aggaatatgg	tgcccgtgtc	660
	tccaacctac	acatgtactc	agctgtctgt	gctgacgtca	tccgccgatg	ggtctaccct	720
	ctcaccocag	aggcgaactt	cactgacagc	accacccaga	gctgcactca	ttcccggcac	780
	aacatctacc	gagggcctga	ggtcagcctg	ggccatggca	gcattcctaga	ggaaaatgtg	840
	ctcctgggct	ctggcactgt	cattggcagc	aattgcttta	tcaccaacag	tgctattggc	900
55	cccggctgcc	acattgggtga	taacgtgggt	ctggaccaga	cctacctgtg	gcagggtgtt	960
	cgagtggcgg	ctggagcaca	gatccatcag	tctctgcttt	gtgacaatgc	tgaggccaag	1020
	atcacagtga	cactgaaacc	acgctctgtc	ctcacttccc	agggtgtcgt	gggcccaaat	1080
	atcacgctgc	ctgagggtctc	ggtgatctct	ttgcaccctc	cagatgcaga	ggaagatgaa	1140
	gatgatggcg	agttcagtga	tgattctggg	gctgaccaag	aaaaggacaa	agtgaagatg	1200
60	aaaggttaca	atccagcaga	agtaggagct	gctggcaagg	gctacctctg	gaaagctgca	1260
	ggcatgaaca	tggaggaaga	ggaggaactg	cagcagaatc	tgtggggact	caagatcaac	1320
	atggaagaag	agagtgaag	tgaaagttag	caaagtatgg	attctgagga	gccggacagc	1380
	cggggagggt	cccctcagat	ggatgacatc	aaagtgttcc	agaatgaagt	tttaggaaca	1440

	ctacagcggg	gcaaagagga	gaacattttct	tgtgacaatc	togtccctgga	aatcaactct	1500
	ctcaagtatg	cctataacgt	aagtctaaag	gaggtgatgc	aggtaactgag	ccacgtggtc	1560
	ctggagttcc	ccctgcaaca	gatggattcc	ccgcttgact	caagccgcta	ctgtgccctg	1620
	ctgcttcctc	tgctaaaggc	ctggagccct	gttttttagga	actacataaa	gcgcgcagcc	1680
5	gaccattttg	aagcgttagc	agccattgag	gacttcttcc	tagagcatga	agctcttggt	1740
	atttccatgg	ccaaggtact	gatggcttcc	taccagctgg	agatccctggc	tgaggaaaca	1800
	attctgagct	ggttcagcca	aagagataca	actgacaagg	gccagcagtt	gcgcaagaat	1860
	caacagctgc	agaggttcat	ccagtggtcta	aaagaggcag	aagaggagtc	atctgaagat	1920
	gactgaagtc	acactgcctg	ctcctttggg	tgtgattgag	tgccctcctg	gctcctgggc	1980
10	tgggacaagt	gaggaactag	ctgcagaggg	atgagtgacc	accatccagg	ctgagactga	2040
	aaggagcaga	ggctggaact	acagtattct	ttcccctgct	agcaaccatg	tgccctccat	2100
	cctgactgtg	gagttgggat	gtggaagtgg	ggctggaaca	aagcttctgc	ctagggagga	2160
	gctaagcagg	cccggcagtt	ggaggaaggc	cagaggaaca	gctttgtgct	ccggctttcc	2220
	ctcagggaac	acgagagagc	agttggctct	ttctgctgct	tgtatatgtt	aatattaaaa	2280
15	gagagtgggtg	t					2291

<210> 10

<211> 1580

<212> DNA

20 <213> Homo sapiens

<400> 10

	atccccctccg	gttttccctca	gtctccacgt	acgtccctca	aagcgcgtcc	taaaaccggg	60
	ataaccggag	cgctcccat	ggaccacacg	gagggcttgc	ccgcggagga	gcgcctgcg	120
25	catgctccat	cgctgggaa	atttggtgag	cgccctccac	ctaaacgact	tactagggaa	180
	gctatgcgaa	attattttaa	agagcgaggg	gatcaaacag	tacttattct	tcatgcaaaa	240
	gttgacacaga	agtcatatgg	aaatgaaaaa	aggttttttt	gcccacctcc	ttgtgtatat	300
	cttatgggca	gcgatggaa	gaaaaaaaaa	gaacaaatgg	aacgcgatgg	ttgtttctgaa	360
	caagagtctc	aaccgtgtgc	atttattggg	ataggaaata	gtgaccaaga	aatgcagcag	420
30	ctaaacttgg	aaggaaaagaa	ctattgcaca	gccaaaacat	tgtatatatc	tgactcagac	480
	aagcgaaagc	acttcatttt	ttctgtaaag	atgttctatg	gcaacagtga	tgacattggg	540
	gtgttcctca	gcaagcggat	aaaagtcac	tccaaacctt	ccaaaaagaa	gcagtcattg	600
	aaaaatgctg	acttatgcat	tgccctcagga	acaaagggtg	ctctgtttaa	tcgactacga	660
	tcccagacag	ttagtaccag	atacttgcat	gtagaaggag	gtaattttca	tgccagttca	720
35	cagcagtggtg	gagccttttt	tattcatctc	ttggatgatg	atgaatcaga	aggagaagaa	780
	ttcacagtcc	gagatgtcta	catccattat	ggacaaacat	gcaaacttgt	gtgctcagtt	840
	actggcatgg	cactcccaag	attgataatt	atgaaagttg	ataagcatac	cgcattattg	900
	gatgcagatg	atcctgtgtc	acaactccat	aaatgtgcat	tttaccttaa	ggatacagaa	960
	agaatgtatt	tgtgcctttc	tcaagaaaga	ataattcaat	ttcaggccac	tccatgtcca	1020
40	aaagaaccac	ataaagagat	gataaatgat	ggcgcttcct	ggacaatcat	tagcacagat	1080
	aaggcagagt	atacatttta	tgagggaatg	ggccctgtcc	ttgccccagt	cactcctgtg	1140
	cctgtggtag	agagccttca	gttgaatggc	ggtggggacg	tagcaatgct	tgaacttaca	1200
	ggacagaatt	tactccaaa	tttacgagtg	tggtttgggg	atgtagaagc	tgaactatg	1260
	tacaggtgtg	gagagagtat	gctctgtgtc	gtcccagaca	tttctgcatt	ccgagaaggt	1320
45	tgagatggg	tccggcaacc	agtccaggtt	ccagtaactt	tgggtccgaaa	tgatggaatc	1380
	atttattcca	ccagccttac	ctttacctac	acaccagaac	cagggccacg	gccacattgc	1440
	agtgtagcag	gagcaatcct	tccagccaat	tcaagccagg	tgccccctaa	cgaatcaaac	1500
	acaaacagcg	agggaagtta	cacaaacgcc	agcacaaatt	caaccagtgt	cacatcatct	1560
50	acagccacag	tggtatccta					1580

<210> 11

<211> 2509

<212> DNA

<213> Homo sapiens

55

<400> 11

	tgggccggggg	atggggcgcc	ggtctgcctt	gacagggttg	caaagttgtt	ttctaaaattc	60
	cgaagcgccc	ctctgcccc	cccccccaat	ctgcttgctg	cggggggtggg	gggtggggggg	120
	gtcacctcct	caggtttcgt	tctttcaaac	tttttgaaac	cctaatttgt	ggcctctgag	180
60	tgggcctcgt	ggactccgc	ctcctaagta	actcttacca	cgtcactagg	ccaaagaggg	240
	gcgtgggggtg	aacgaaaggg	ctcccgaact	tttttttttc	cagccaggcc	gaacggggggc	300
	tcggtaatga	ttggccaggg	cgcatacactg	cgaacctgtc	aatcacgggt	cctccgggtt	360
	gcgagggggcg	gaccaagccc	caaccccggg	gaatccgagc	aggtatataa	ggggcccagc	420

	tagagcccag	gcagactgtg	aatgcgacct	gttcgagaga	actcatcagg	tgcgagaagc	480
	ccgcgggttc	ctgctgattt	ggcgcgagc	attttgataa	gcctaccctt	cccgcgggac	540
	tcgctggccc	acagggcccc	aagctccgct	ccgacggagt	cccagggcct	tttcaccgtg	600
	gcccgtccag	ccccgggagc	gccttctcct	cccgccacgc	tggcgcaact	tcttcccgcc	660
5	ccggcaatgt	acagccttct	ggagactgaa	ctcaagaacc	ccgtagggac	accacacaaa	720
	gcgggcgggca	ccggcgggcc	cgcagccccg	ggaggcgag	gcaagagtag	tgcgaacgca	780
	gcccggcgcg	cgaactcggg	cggcggcagc	agcggtggtg	cgagcggagg	tggcgggggg	840
	acagaccagg	accgtgtgaa	acggccccatg	aacgccttca	tggtaggtgc	ccgcggggcag	900
	cggcgcaaaa	tggccctgga	gaacccccaa	atgcacaatt	ctgagatcag	caagcgcttg	960
10	ggcgccgact	ggaaactgct	gaccgacgcc	gagaagcgac	cattcatcga	cgaggccaag	1020
	cgacttcgcg	ccgtgcacat	gaaggagtat	ccggactaca	agtaccgacc	gcgcgcgaag	1080
	accaagacgc	tgctcaagaa	agataagtac	tccctgcccc	gcggcctcct	gcctcccggt	1140
	gccgcggccc	ccgcgcgcgc	tgccgcggcc	gcagccgctg	ccgccagcag	tccggtgggc	1200
	gtggggccagc	gcctggacac	gtacacgcac	gtgaacggct	gggccaacgc	cgcgtagctg	1260
15	ctgggtgcagg	agcagctggg	ctacgcgcag	cccccgagca	tgagcagccc	gcgcgcgcgc	1320
	cccgcgctgc	accgctacga	catggccggc	ctgcagtaca	gccccaatgat	gcgcgcgcgc	1380
	gctcagagct	acatgaacgt	cgctgcgcgc	gccgcgcgcg	cctcgggcta	cgggggcatg	1440
	gcgcctcag	ccacagcagc	cgcgcccgcc	gcctacgggc	agcagcccgc	caccgcgcgc	1500
	gcgcgagctg	cggccgcagc	cgccatgagc	ctggggcccca	tgggctcggt	agtgaagtct	1560
20	gagcccagct	cgccgcgcgc	cgccatcgca	tcgcactctc	agcgcgcgtg	cctcggcgac	1620
	ctgcgcgaca	tgatcagcat	gtacctgcca	cccgcggggg	acgcggccga	cgccgcctct	1680
	ccgctgcccc	gcggtgcgct	gcacggcggtg	caccagcact	accagggcgc	cgggactgca	1740
	gtcaacggaa	cgggtgcgct	gacccacatc	tgagcaccgc	cctgcgctcg	tccacccttg	1800
	ttccccaccc	ccacccccac	tcccgccccg	caccccccaag	ttgggtcgcc	ttgttttagct	1860
25	ttgcttgctt	gggactgttg	ccttgtagcc	atgatgggga	gggctgaaag	ttttgctgta	1920
	gctgtcggtt	tttgtacaaa	agtcaaaaat	aagtcaggag	cagcgaaaat	gggatcttct	1980
	agagagctct	cttgccccac	gccgctgctc	ctttcacctt	tgtaggctgg	gaatcgctgt	2040
	gttatttgca	aagaaaaaac	agccccact	cctcctcctg	agttccaggg	ttattctggt	2100
	acatttgaaa	atgttgtctt	gttagtttgc	agtttagcaa	ggagtgaatg	ggagaaacat	2160
30	agtatcggtt	gaggtccagc	tggagaactg	caacgcctac	gccccagtc	gtgtcgctc	2220
	tgttttcctc	caggtttttt	ggggcgctga	ccgctccaag	cagcgcgcca	gctaaagcca	2280
	atgttaattt	atagccagg	gtgcgtgtgt	ctccgcctc	gccgcccctg	gccgcgggac	2340
	agcttctgtc	caatcatgtt	gagttgggtg	tttctgccgt	gatctgtttg	atattttctt	2400
	gcgctaattg	gttcagattt	cgtttgggta	gtggggaggg	gctaactttg	ttcagggttt	2460
35	tcaagctttt	actcttaatt	cctaaatgag	atcaataaat	tttataacc		2509

<210> 12

<211> 8372

<212> DNA

40 <213> Homo sapiens

<400> 12

	aagcttgggtg	ccatctattt	tggaactatgc	cttgcataca	gcttttatggg	aacatttgtc	60
	aggcaaaaagt	ataataatgg	caaactctac	gccttttatt	ttaaattaga	ttgggtgtgat	120
45	ttgatgctga	cgggagtgag	agtaatggcc	ttatcctgct	gcaggctgtg	ctgaggatgg	180
	cctggtctgc	caccctcctc	gagtagcatt	ttgcatgtgt	aacagggtct	cccctctggg	240
	gcacaacaac	aaagagaagt	tgctaaggac	aagaagcagg	tgcggaaatg	catctcccat	300
	tggaaacagc	ctgggcttac	tccaatggct	gagagaggtg	ctatggccag	tcctcccaga	360
	gctctgcagc	tgcacttggg	ggtggacagt	ctcgtgcttg	tcctgcgtga	taacggccgt	420
50	gaaagccagc	caactgctgc	ccaaaatcac	ccagccgatt	gggggtttcc	catcggcgca	480
	ccctgcccgg	agccaagaag	acaggctggt	gctgctgtat	ttgtatttat	atccattgct	540
	gcgctctgcg	ttctcgtggc	acgcctggac	actcctccgc	ctccccctcc	tcttccctct	600
	ccagggccac	ctccccgcct	tccccacccc	catctgcttc	tgtcaaatga	gaaagtcacc	660
	gaggagaacc	caaacactcc	agccgctgag	agcccccttt	ggcacttggc	agcacgcggc	720
55	ggcgggctcc	tcggctcaac	ttcgaggagt	ctccgcgacg	caacttttgg	ggacgctttg	780
	catttaagag	agaacgaccg	aggaggagga	gcgctctgcc	cggccgcgcg	tacctgcggg	840
	gagctcacca	gcaaacgcca	ctgcagacga	aggacccaaa	gaacgtaaag	ggcaaacctg	900
	cgccgcgggg	aggggggcacc	gccgagaagt	tagagtgtcc	cagagacaac	ctgctcgagc	960
	gctcggcggg	agacactaag	ggggcccggg	gcgcggcggtg	gccctggctg	gtcccccagc	1020
60	cccctcctcc	ggggcgggag	cgacgcgggg	gcgcgacgag	ccccggccgg	ccgagcgggt	1080
	ctccgcgggg	agccaacatt	gatttcctcc	gggcccaggg	cgagggcccg	ggcgcgggcg	1140
	ggctgcagcc	gcggcgaggc	gagagcatgt	ccaagccggt	ggaccacgtc	aagcgcccca	1200
	tgaacgcctt	catggtgtgg	tcgcggggctc	agcggcgcaa	gatggcccag	gagaacccca	1260

	agatgcacaa	ctcggagatc	agcaagcgct	tgggggcgca	gtggaaactg	ctcacagagt	1320
	cggagaagcg	gccgttcac	gacgaggcca	agcgtctacg	cgccatgcac	atgaaggagc	1380
	accccgaact	caagtaccgg	ccgcggcgca	agcccaagac	gctcctcaag	aaggacaagt	1440
	tcgccttccc	gggtgccctac	ggcctgggcg	gcgtggcgga	cgccgagcac	cctgcgctca	1500
5	aggcggggcg	ggggctgcac	gcggggggcg	gcggggcgct	ggtgcctgag	tcgctgctcg	1560
	ccaatcccga	gaaggcgggc	gcggcccgcg	ccgctgcgcg	cgcacgcgct	ttcttcccgc	1620
	agtcggccgc	tgccgcggcc	gctgcccgcg	ccgcgcgcgc	cgcgggcagc	ccctactcgc	1680
	tgctcgacct	gggctccaaa	atggcagaga	tctcgctgct	ctcgtccggc	ctcccgtacg	1740
	cgctcgctcg	gggctaccgg	accgcggggc	cggggcgcct	ccacggcgcg	gcggcgggcg	1800
10	ctgcagcggc	ggccgcggcc	gccggggggc	acacgcactc	gcaccccagc	ccgggcaacc	1860
	cgggctacat	gatcccgtgc	aactgcagcg	cgtggcccag	ccccgggctg	cagccgcgcg	1920
	tcgcctacat	cctgctgccc	ggcatggggc	agccccagct	ggacccctac	cccgcggcct	1980
	acgctgcccg	gctatgacct	cgcggggccg	cctgcgcagg	accggtgtgc	acacgtgtac	2040
	atatgtatag	gtacgagcgc	tgccgcctcc	ccgtgcgcgc	tcccgcgacc	ggggggcccg	2100
15	tttgtatgta	catagaatgt	ataggtgcc	ggtagaggca	gagaggccag	gcggggcagg	2160
	agtggccaag	cgcgcaaggg	cgcggggcag	caggcctgtg	aattcgcagg	atcatttcag	2220
	acccgcactt	cggcagccaa	ctcgaaagca	ggcgggtgtg	tgccgcagca	gttggcgttt	2280
	gctttgcact	tcggaacctg	ttgcgttttg	accacgggag	gtggaggagt	aactttttga	2340
	catgtttggc	tttccagttt	tggtggaagt	ttcatggtcg	gtttttgttt	tgttttctcat	2400
20	tcttcttccc	cgcccccag	cccccaacc	cccaaccccc	tcccggtcgc	tggtgcagtc	2460
	acgctgttca	aatgtgaggt	ctgaaatggc	tgccacacgc	gaaaagctgc	ttgtgtcatt	2520
	cgtttctggg	agtgggatgg	ctctgagcag	cctgcgcctc	ctgtttgtac	tatttgaact	2580
	ttgcagatct	ctgttctctc	aagcagaact	cccaaccaga	tccattcttg	accagtacc	2640
	ggctcgaatc	tggccttttg	tgtgagatga	tcacggnntc	ttttgtttat	caagccattt	2700
25	gcaaatcaga	gcaagagctc	tttctcaagg	gcaagaaacg	caaacaagaa	atattttgtga	2760
	gatgaaagtt	gtcaattgga	tttcttccc	aaacaacaaa	caacaacaaa	ctactagaag	2820
	tctccctgag	tccactcgct	tggatttctg	acacagttta	caaaaaagga	aaaaggcact	2880
	gctcctattt	tcccttatgg	ctgagttcac	cttaagattg	taaatgtgta	tatgtcagtg	2940
	aaaacattga	ggccttgaaa	atgtgttatt	ttcgttgccc	taagtttgag	tcgactttag	3000
30	actcaaaaac	attttgagcg	aatatcaaag	ttacttttta	aaaattgcga	aactatttca	3060
	gaatcgcaat	tttatcgaag	attaaatcag	acttttttgt	ctggttaatta	tatatattat	3120
	atttagcaaa	actgaagaaa	aaaagcacag	aattgtttca	acagatgtct	ctcattttca	3180
	gctagcattt	ctctcccaag	ttgagctggt	ttaatgtggt	ttggatttcc	ctcctcaatt	3240
	ggcttatttt	ttagatcacc	tgcaattcat	ttgcaaattg	caataaaaca	catttttagaa	3300
35	aaaaggaacc	ttcaattatt	agctttgttt	ctttttaaat	gtatatat	tgactaatgt	3360
	ttgtgaatga	agttggctaa	catgtattta	gtttcatttt	ggctttatgt	aatataaagt	3420
	ttttaaaatt	ttaaatatgg	ttttaacctt	tatgtgtaaa	tgattttcta	gtgtgacctt	3480
	ctaattttaa	attagacgtc	taaggtatat	ctgtaaaatta	gaatccgact	atcactctgt	3540
	tcattttttt	tgaacaaaga	gtttaaataa	agcctgaacc	agggaaaaga	aaaatcttct	3600
40	atttcttgtt	gagttcctaa	caagattttt	atctgaattg	cccttacgtg	cctgggtccag	3660
	gtgaagtgtg	aggtatcctc	caaaggcacc	ctttgtttca	cttttgaata	gatttactag	3720
	gaaatctaaa	tcaagccatt	gttattcaga	gccaaaaacc	tgatttatca	cattttttaa	3780
	cgtgaatagg	aaagaagatt	tttaaaaagc	ccaagtcgtt	gtattagctt	taacaacaac	3840
	aaaaaaaaag	cattcatgaa	ccagtagaac	agagcccatt	gaaaacatcc	agacctttca	3900
45	aagcatttca	ccagtttcta	gtaacatttt	aagaggggaa	agttgcttga	ccactttatc	3960
	ttgttagttg	aagagcccca	ccacttaaat	cagtgtaat	tggtctccta	tctttggggg	4020
	attccttgtt	gacaccttaa	ggtttttatt	ggaaggataa	tcactactaa	cgacaaagta	4080
	caaatttttg	cctcttttag	acttaatttt	gttatgctaa	tcgcattaaa	gtagaagtat	4140
	aacattcaaa	tggagagggg	tggatttcta	gggctagaca	aattgctact	aaagtttgaa	4200
50	aaatcataaa	gggattttat	tttagacaag	aaatagaaga	ctgtcagaaa	aaaaaaaaata	4260
	ggaagatctc	gccccccgcg	aacaaaaatg	gaaattctca	agatactata	tacaagtctt	4320
	aaaccagttt	ccccattgag	accatctctg	gagctgcacg	tctttataaa	cgacccaagt	4380
	ctttaaagtc	attgttttcc	cccaacggaa	taatatttta	aaaaccatga	aaagttttgg	4440
	aaatgtgaga	aataggctct	gctgggttga	ccctgattca	ctaattaaaa	tgatccctct	4500
55	cctgttatcc	cctgagctct	ttgcaatatt	ataagttaat	tcatatgggt	ctgagcgatt	4560
	atgcaaaaac	aatttgagct	gtccaggggt	aattatccct	gacacgggta	attaaatcct	4620
	ttcaaggcct	ogtctttccc	ttttgtagca	gcccatacct	tctcaacacg	gaacttctgc	4680
	ggctcgctgg	aaatcacccc	agccctaaat	ccttagttacc	accctgagcc	ttccagctcg	4740
	gccgcctcct	cggcctgaag	actccccgcc	tcctcccgcc	ccctcccttt	ttcccaaaga	4800
60	tcagcggttt	ctgggagaaa	cgcctccggg	ttgttgatga	atgagaagag	gactggaaaag	4860
	atgggtaaga	ggaggggtga	ggatgcogag	ggggagcacc	gaggtcatat	cgccaacaga	4920
	ttgtgcggct	gtttgaggac	ctccacaggg	cccacagact	cgtttatcac	ccattctgac	4980
	tccaatggct	ttgctaacaa	ggtggcgggg	tttgcgcctg	cagagagcct	cctgccaaagt	5040

	tagactgtgc	agaagtaagg	ggttgagcgc	gggggagcgc	ctccggggca	agagggcgta	5100
	gagaaaggcc	cggggnnggg	nggtgtaagc	gtctgaaagt	ggcccacaaa	tgcagcgctg	5160
	tgattgggca	gagagctgct	gctggctcgc	gatctctatc	tccatctctt	tatctatctc	5220
	cgtctctctc	cctgtttctc	catttttctt	tctttccctt	tctctccttc	cttccctcca	5280
5	tctttctctt	ttcccttcct	tttattcttc	tattttcggt	tcttttcaag	gtttttttta	5340
	aagccatgat	gcaatttctt	tggtattcac	cgttggtcca	aaaottgaag	caagcctcgt	5400
	atccaagggg	ccaggcatgt	tgcttcgggc	tttgtgcaaa	caggtggaat	tgcgctgtgt	5460
	aagcagtaag	aactggtgct	ggggagctgt	cgcgcgaggg	ggtggctttg	ggagagcagg	5520
	gttgctggcc	gcgattgtta	cttcccttga	caatttcctc	ctccccctcc	cccaagaaga	5580
10	taggagaaag	caccgcggat	ctccctctca	ccccaggctc	ggggcgcgaga	agatggagag	5640
	aagattccac	tctccccgga	gcagataggg	acggctcgcg	cagccaatca	gagcgcggt	5700
	cggcgccggc	gctccccggc	gcctgggccc	ccgtgtcctc	caggcaagcg	aagtccccgc	5760
	aactcgtccg	cctcgagggt	cgcgctcttt	cttgcccccg	cggcccagcg	gaggccgagg	5820
	gagccgtcca	aactttatta	atctctcctc	ctttctttct	ccctcagccc	agtgcattctc	5880
15	aaaggtcagc	cctcttcttt	taaaagactg	atattattaa	tgcactgaca	attcctcccc	5940
	cccttttctt	ttttctctct	tgaggggggg	aaaaaaaggg	aaatggtgaa	aagagctttt	6000
	tttatccttt	tttttttttt	gtccttcagt	gggagcgttt	agacagtcga	ggagggtttg	6060
	tccgagaaca	aaacgcaggg	ttgggaggtt	ttgtgagagt	gttggtttgt	gaagtggagc	6120
	taagaaaaag	cggcggtctt	ctcctcattg	tgaagaaaac	aatcagtggt	atltggaaaa	6180
20	ctgttagcat	tgtgcacttc	ttctgtgtcc	attgtgaggc	gtttcttttc	acaagggttt	6240
	tttttcagcc	gatccagctg	gccggaatga	atagcgtgtc	aatgtgtaca	cgctttgtcc	6300
	ctccggcctt	caagttagcc	ccattgaata	gactaagttg	acctgcgtga	cagtgaacaa	6360
	acataataaa	aaatacatga	gcccctgaat	aggagcaggg	gcataaataa	ataaaatggg	6420
	tgacaaaaac	tggataaact	gaatgacaaa	acggtgaaaag	gggaacaaaa	agatatattaa	6480
25	cacgctagat	tagcattaga	atgcgatcta	caaggcagaa	caattgatga	atagggtttac	6540
	cggccaagaa	agaaatggac	taaatgccct	ttgaatagat	atgctttttg	caagggtctt	6600
	gaatagatat	gcttttgcaa	gggctgaatg	ggaaaaggta	aagatgaagc	tatgcaaattg	6660
	agccggggaa	ctttttatat	atattcttta	aacacacaca	cacactgcgg	ggggaagagt	6720
	gctgcctcgg	gatgtttata	gaagcaataa	ttgccattat	tagcattgtc	tgcggcagat	6780
30	agaaattgaa	cagggttgga	taatataagg	tagcagtaat	tattcttcta	attaatgggtc	6840
	ctttgtctact	tgaaaaaaga	aaaaaggaaa	gaagttagtaa	aagttatgca	gaagttatgt	6900
	ttccttgtgt	ccatttgccc	agcgttgaa	tctgtggagc	aggaagcctg	gcaattccaa	6960
	gatacgcgat	gatcytcaaa	cattccccgg	agccagtcct	gaggtctctg	cttcaggggc	7020
	tagttttccat	ttatgccgcg	tttttgagag	tctaatactg	tgtctggcac	atggtaggtg	7080
35	ctcactgaat	agtcgtggta	tgaatgaatg	aacgaatgaa	tgaatgaatg	aatgaatata	7140
	agtttaaatg	gggaaacccg	ggcctcctaa	taaaggtagg	ggctggggga	tacctagggg	7200
	cttccccagg	aggatttctt	ttttcatcat	cccacccctg	ggagaaagg	ccacgcagga	7260
	tggtegtctc	ccccttgctg	agagttttgc	cttcagccta	tctgggcccgc	tggaaaagag	7320
	gagaaagata	aacaagagac	aagcaactac	tcccctaccg	gcgttccgtc	cttgctctca	7380
40	ctgccaaatc	cactccaaag	ccgaggatgg	tgagactgtg	aagttgcaaa	gaaacacaga	7440
	gcccaccccc	ttaaagaatt	acgatataat	taaagtttgc	ctctttcagg	tttctctcct	7500
	tggctcctgc	ccctttcccc	tcccggctcc	ttgtccttga	ctgaacctca	tgggacagag	7560
	aacctcctgt	cccccacgag	gcaaggcgcg	aaccgcgaga	gatctggggg	gccctttggt	7620
	tccctgcgct	gccctggagg	cgtccataga	ggcctttgcc	gccaaggaca	gcaattgttt	7680
45	tattttcgat	ggttgctcgc	caggctgcgg	gtcgcggggc	caccagcccg	tcgaaactttc	7740
	cagtcgttat	cagcgtgtgt	cctaacttaa	tggaaataatg	caaattatag	cctgcccagc	7800
	tgacacgtcc	ctgcgaatgc	gccggggctg	agctctggcc	agccgtctct	tgcagctcct	7860
	ggacggccgg	agggaatgaa	gctctgaatt	gtgacaaaag	tggggggggc	accccaaatt	7920
	ctcaaagcaa	tgttcttttt	tttttctttt	ttcttaagca	attgagcctt	accaaattgtc	7980
50	ggggccggcc	gcaagggaag	cttgcatatt	ttaaagtgtg	acctgagcct	tgcgggtttc	8040
	agcttcaact	aaaacatgca	aattcttgaa	attgaaaaat	ctgaaaaact	tccgaagagt	8100
	tctatctgaa	taaattccaaa	tccattggga	gtcgttttga	ggagacaaaa	cgcacagcga	8160
	tttggggtga	gggatatttt	tggggaggca	ggacgtgctg	gattgggttt	ccagggtcaa	8220
	ggtgtctctg	ggccttcgac	gatagcctta	gcgcagagca	gggaagtggc	accgctaggc	8280
55	agcaagctca	gttgtctctc	ttttgtgacc	catcccccca	cccccccccac	cgcacacctt	8340
	gctccggggc	cactgccccct	ctctgcaagc	tt			8372

<210> 13  
 <211> 4877  
 <212> DNA  
 <213> Homo sapiens

<400> 13

	gcccga	aacc	cggaagt	gag	cggcggc	gagc	tgcgagg	ctc	ggagaa	acag	gcgccgc	ggg	60
	ctccgc	gcccc	ggccgg	accc	gggcccc	gaga	tcatgat	gct	gccgcc	caccg	ccgccacc	ac	120
	ggagcg	agaa	ccccag	atag	acgcccc	ggc	ggcccc	gggt	cctggag	tcc	cgccgc	cctgc	180
	tgcccc	ggccg	aggac	ccccac	cccgcc	tgcc	gcccgat	gct	tgcagt	gggg	cccgcc	atgg	240
5	acaggg	atta	cccgca	gcac	gaaccccc	gcg	cggcggg	cag	cctcct	gtac	agcccc	gcgc	300
	ccctgc	agag	cgccat	gctg	cactgccc	ct	actgga	acac	cttctc	gctg	ccgccata	acc	360
	ctgcct	ttctc	cagcgac	agc	cgcccc	gttca	tgagct	ccgc	ctcctt	ctctc	ggcagcc	agc	420
	cctgccc	caga	caccag	ctat	gcccccg	tgg	ccaccgc	cctc	cagctt	tgcca	ccaaag	acct	480
	gcgact	tttgc	tcagg	actcc	tcctat	ttttg	aggact	ttctc	caacat	ctcc	atcttct	cct	540
10	cgtcct	gtgga	ctccct	gtcg	gacatc	gtgg	acacgccc	ga	cttctc	gtccg	gctgac	agcc	600
	tcaacc	agg	gtccacc	atc	tgggac	gata	acctgccc	c	ctccacc	ccac	gataag	ctgt	660
	tccagct	cag	caggcc	gttt	gcaggct	tctc	aggact	ttctc	gccctc	ccac	agcacc	ccgc	720
	ttctcgt	cag	ctacc	aggag	cagagt	gtgc	agagcc	agcc	agagg	aggag	gacgag	gctg	780
	aggagg	aggga	ggcgg	aggag	ctgggg	caca	cagagac	ctc	cgccga	ctac	gtgccg	tcca	840
15	agtcca	agat	cggaag	cag	caccag	acc	gcgtgg	tgga	gaccag	caca	ctgtcc	agcg	900
	tcccc	cccc	agacat	cacc	tacacct	tgg	ccctgc	cctc	ggacag	cggg	gccctg	tctg	960
	ccctgc	agct	agagg	ccatc	acctac	gcct	gccagca	aca	cgaggt	ctctg	ctcccc	agcg	1020
	ggcagc	gcgc	gggctt	ttctc	atcggc	gatg	gggccg	gct	gggcaa	aggc	cggacg	gtgg	1080
	ccggag	tcat	cctgg	gagaac	cacctg	cgcg	gccgga	agaa	agcatt	gtgg	ttcagc	gtct	1140
20	ccaacg	acct	caagt	acgat	gcgga	gcgc	acctgc	ggga	catcga	agcc	acgggc	atcg	1200
	cggtgc	acgc	gctcag	caag	atcaag	tacg	gtgacac	cac	tacct	cagag	ggcgtc	ctct	1260
	tgccac	ctta	ctccg	ccctg	attggg	gaga	ccaggcc	cg	tggcc	agcac	cgcact	gcgc	1320
	tccggc	agat	cctgg	actgg	tgtggg	gagg	cctttg	aggg	cgtcat	cg	ttcgac	gagt	1380
25	gtcaca	aaagc	caaga	atgcc	ggctcc	acca	agatgg	gcaa	ggccgt	gtcta	gacctg	caga	1440
	acaagc	tgcc	cctgg	ccccgc	gtggtc	tacg	ccagcg	ccac	aggtgc	ctct	gagcct	cgga	1500
	acatga	tcta	catgag	ccgc	ttgggt	tatct	ggggcg	agg	cacacc	cttc	cggaac	tttg	1560
	aggagt	tcct	gcacgc	catc	gagaag	agg	gcgttg	gcgc	catgga	gatc	gtggcc	atgg	1620
	acatga	agg	cagcgg	cag	tacatc	gcac	gccagc	t	cttctc	ccggc	gtcac	cttcc	1680
	gcacgc	agga	gatccc	gcgt	gcccc	agcct	tcgagt	gcgt	ctaca	aaccgc	gcagcc	ctgc	1740
30	tgtggg	ccga	ggccc	tgaac	gtgttc	cage	aggcgg	ccga	ctggat	cggc	ctggag	tgcg	1800
	gcaagt	ccct	gtggg	ccag	ttctgt	tcgg	cacacc	agc	cttctt	caag	tatctg	tgca	1860
	tgcagc	caa	ggtgc	gcgc	ctggtg	gagc	tggccc	gaga	ggagct	ggcg	cgagac	aaagt	1920
	gcgtgg	tcat	cgggt	gcag	tcacg	gggc	aggcgc	gcac	gcggg	agg	ctggg	gggaga	1980
	acgatg	ggga	cctca	actgc	ttcgt	tctcg	ccgtga	agg	cgtgtt	ctctg	tcgcta	aattc	2040
35	agaagc	actt	tcctg	tccac	aagaga	aaagc	gggac	agag	agcggg	cagc	aagcgg	aaac	2100
	ggcgac	ctcg	gggac	gcggg	gccaa	agccc	cccgg	ctggc	gtgcga	agaca	gcggg	cg	2160
	tccgc	atcag	tgacg	acagc	agcag	cgag	cggacc	ctgg	cctgg	acagc	gacttc	aaact	2220
	cctcccc	cga	gtccct	ggtg	gatgac	gacg	ttgtc	atcgt	tgatg	cagtc	gggctc	ccca	2280
	gtgacg	acgg	gggat	ccctg	tgcctc	ctgc	agagag	acc	gcac	ggcccc	ggggtc	ctgg	2340
40	agcggg	tggga	gcgggt	gga	caggat	ctgc	tggaca	aaagt	gcgccg	gctg	ggccgg	gaac	2400
	tgccag	tcaa	cacct	ggac	gagctc	atcg	accag	ctggg	cgcccc	ccag	cggg	gtggcg	2460
	agatga	ccgg	caggaa	aggc	cgctg	gtgt	ccagg	ccccga	cggg	acggtg	gcctt	cgagt	2520
	cgcggg	caga	gcagg	gtctg	tccat	cgacc	acgtga	acct	caggga	gaga	cagcgt	ttca	2580
	tgagc	ggcga	gaagc	tcgtg	gccat	catct	cggag	gcctc	cagctc	gggt	gtctcc	ctcc	2640
45	aagccg	accg	ccgtg	tccag	aaccag	cggc	gccgc	gtgca	catgac	cttg	gagctg	gccgt	2700
	ggagcg	ccga	ccgcgc	catc	cagcag	ttcg	gccgc	accca	ccggtc	ccaac	caggtc	ctccg	2760
	cgccag	agta	tgtctt	ctctc	atctcg	gagc	tggccg	ggga	gcgccg	gttc	gcctcc	atcg	2820
	tggcca	agcg	cctgg	agagt	ctgggg	gccc	tgacc	acag	agacc	gcgc	gccacg	gagt	2880
	cccgtg	acct	cagca	agtac	aacttt	gaga	acaagt	atgg	cacc	cgggc	ctgca	ctgtg	2940
50	tcctca	ccac	catcct	gagc	cagact	gaga	acaa	agtgcc	tgtg	ccccag	ggata	ccctg	3000
	gagggg	tccc	cacctt	cttctc	cgggac	atga	agcagg	gcct	gctgt	ctgtg	ggcatt	gggtg	3060
	gccggg	agtc	ccgga	atggc	tgcctg	gacg	tggaga	agga	ctgtt	ccatc	accaag	ttcc	3120
	tgaacc	gcac	cctggg	gctg	gaggtg	caca	agcaga	aatgc	cctgtt	ccag	tactt	ctcag	3180
	acacct	tcga	ccacct	catc	gagatg	ggaca	agcggg	agg	caaata	cagac	atggg	catcc	3240
55	tggacct	ttgc	tccgg	tatc	gaggag	atct	acgagg	agag	ccagc	aggtg	ttcctg	ggctc	3300
	ccgggc	accc	caggac	cg	caggtg	gtct	ctaca	agat	cagcgt	ggag	cgcgcg	ctga	3360
	agtggg	aggga	cgctt	ttgcc	aagtc	ctgtg	cctgac	cg	cccct	atgac	ggctt	ctacc	3420
	ttctct	ataaa	ggtcc	gcgg	gt	aaca	agcccc	a	gctgc	ctgct	ggcgg	agcag	3480
	agttct	ttcac	ggtgt	tacaag	cccaac	atcg	gccgg	cagag	ccagc	tg	ggag	agtg	3540
60	gcctcc	gcgc	caagt	ttccac	cggtt	caccg	cggag	agggc	caagg	agccc	tggg	agagtg	3600
	gctacg	cttt	gtcgt	gacg	cactg	cagcc	acagc	gcctg	gaacc	ggcac	tgccg	gctg	3660
	cgcagg	agg	taagg	actgc	ctgc	aggggc	tgcg	gctgc	gcacc	actac	atgct	gtgcg	3720
	gcgcgt	gct	gcgcgt	gtgg	ggccg	catcg	ccgc	cg	ctc	at	agcag	cagca	3780

	gctacctgca	gatcgtgagg	ctgaagacca	aggacaggaa	gaagcaagtg	ggcatcaaga	3840
	tccccgaggg	ctgctgtgag	cgggtgctgc	aggagctgag	gctgatggat	gcggacgtga	3900
	agcgcaggga	ggcgcccgcc	ctgggtgctgc	cgcgcccgcc	cgcccccgcc	ccgctggcgc	3960
	tgccttgagg	ccccggagag	gtgctggacc	tcacctacag	ccccccggcc	gaggccttcc	4020
5	cgcgcgcccc	gcacttctct	ttccccgggc	cgtgttccct	ggacgcgggc	cccggtctcg	4080
	tgcgctggg	cacccccgac	gcccaggccg	accctgcggc	cctcgcgcac	cagggctgcg	4140
	acatcaactt	caaggagggtg	ctggaggaca	tgtctgcctc	gctgcacgcg	gggcccgcct	4200
	ccgagggcgc	gctgggggag	ggcgcggggg	cggggggcgc	ggcgggcggt	ggtcccgcgc	4260
	ggcagagcgt	gatccagttc	agcccaccct	tccccggcgc	ccaggctcct	ctctgacacg	4320
10	ccttttaggg	aaacatgccc	caagacacag	ggaccgtttc	tccccctagga	gcagcgggtg	4380
	ggagcagggc	caagggtcccc	tgaccactgc	tcagaggagc	cctaggccct	ggccgcagtg	4440
	ccttcagcgc	ccgacccggg	ccccacactg	gtcagccctg	gcggggccca	ctcaggacag	4500
	ctggggggcg	gggctgtgga	gggcccctct	tgtgcctctc	ctcctaagta	ggaaggggct	4560
	ccgggtggct	gctctgggac	tgggcaccca	caagggtcca	gtggggccaa	acccttgaaa	4620
15	tccgtgaaac	cgggtgggtc	caagagctag	aaactcagga	aaccccaggt	gctcagggcc	4680
	ccgcgtctcg	ggggctccgt	ggggcagacc	cctgctaata	tatgcaattc	tccctcccc	4740
	agcccttccc	tgacccctaa	gttattgccc	gctcacctct	cccaggcccc	aggccgcgga	4800
	gctggcaggg	tggcgccctg	ggtttctatg	tatttatagc	aagttctgat	gtacatatgt	4860
	aaaggacttt	tttaaat					4877
20	<210> 14						
	<211> 1872						
	<212> DNA						
	<213> Homo sapiens						
25	<400> 14						
	tcaggctgcc	tgatctgccc	agctttccag	ctttcctctg	gattccggcc	tctggtcata	60
	cctccccacc	ctctctccaa	ggccctctcc	tggctctccct	tcttctagaa	cccccttccc	120
	cacctccctc	tctgcagaac	ttctccttta	ccccccaccc	cccaccactg	ccccctttcc	180
30	ttttctgaac	tccttttgga	gggtctcagc	ctgccagac	cataggagag	atgtgggagg	240
	ctcagttcct	gggcttgctg	tttctgcagc	cgttttgggt	ggctccagtg	aagcctctcc	300
	agccaggggc	tgaggtccc	gtggtgtggg	cccaggaggg	ggctcctgcc	cagctccctc	360
	gcagcccccac	aatccccctc	caggatctca	gccttctgcg	aagagcaggg	gtcacttggc	420
	agcatcagcc	agacagtggc	ccgcccgtcg	ccgcccccg	ccatccctcg	gcccccgcc	480
35	ctcaccgggc	ggcgccctcc	tcctgggggc	ccaggccccg	ccgctacacg	gtgctgagcg	540
	tgggtcccg	aggcctgcgc	agcgggaggc	tgcccctgca	gccccgcgtc	cagctggatg	600
	agcgcggcgc	gcagcgcggg	gacttctcgc	tatggctgcg	cccagcccg	cgcgcggacg	660
	ccggcgagta	ccgcgcgcgc	gtgcacctca	gggaccgcgc	cctctcctgc	cgcctccgtc	720
	tgcgcctggg	ccaggcctcg	atgactgcca	gccccccagg	atctctcaga	gcctccgact	780
40	gggtcatttt	gaactgtctc	ttcagccgcc	ctgaccggcc	agcctctgtg	cattgggtcc	840
	ggaaaccggg	ccaggggccga	gtccctgtcc	gggagtcccc	ccatcaccac	ttagcggaaa	900
	gcttcctctt	cctgccccaa	gtcagcccca	tggactctgg	gcccctgggg	tgcatcctca	960
	cctacagaga	tggcttcaac	gtctccatca	tgtataacct	cactgttctg	ggtctggagc	1020
	ccccaaactc	cttgacagtg	tacgctggag	caggttccag	ggtggggctg	ccctgccgcc	1080
45	tgcctgctgg	tgtggggacc	cggcttttcc	tcactgccaa	gtggactcct	cctgggggag	1140
	gccctgacct	cctggtgact	ggagacaatg	gcgactttac	ccttcgacta	gaggatgtga	1200
	gccaggccca	ggctgggacc	tacacctgcc	atatccatct	gcaggaacag	cagctcaatg	1260
	ccactgtcac	attggcaatc	atcacagtga	ctcccaaata	ctttgggtca	cctggatccc	1320
	tggggaagct	gctttgtgag	gtgactccag	tatctggaca	agaacgcttt	gtgtggagct	1380
50	ctctggacac	cccatcccag	aggagtttct	caggaccttg	gctggaggca	caggaggccc	1440
	agctcctttc	ccagccttgg	caatgccagc	tgtaccaggg	ggagaggctt	cttggagcag	1500
	cagtgtactt	cacagagctg	tctagcccag	gtgcccacag	ctctgggaga	gccccagggt	1560
	ccctcccagc	aggccacctc	ctgctgtttc	tcaccttggt	tgtcctttct	ctgctccttt	1620
	tgggtgactgg	agccttttgg	tttcaccttt	ggagaagaca	gtggcgacca	agacgatttt	1680
55	ctgccttaga	gcaagggtat	cacctctgcc	aggctcagag	caagatagag	gagctggagc	1740
	aagaaccgga	gccggagccg	gagccggaac	cggagcccga	gcccagagcc	gagccggagc	1800
	agctctgacc	tggagctgag	gcagccagca	gatctcagca	gcccagttca	aataaacgtc	1860
	ctgtctagca	gc					1872
60	<210> 15						
	<211> 1201						
	<212> DNA						
	<213> Homo sapiens						

<400> 15

	gagtctacgg	cattgctgag	gacgctgccc	agggcatcgc	taatgaggac	gccgaccagg	60
	gcacgcgttaa	tgaggacacc	accagtgca	tcgccaacga	ggaagccgcc	cagggcatcg	120
5	ccgaggacgc	catccagggc	atcgccaacg	aggaggttgc	ccagggcatc	gccaatgggg	180
	tcgccgcaca	gggcatcgcc	aatgaggacg	ccaccagggg	catcgccaac	tgggacgccg	240
	tcacggcctt	cgccaacggg	gacgccgtcc	tcagcttcgc	caacggggac	gccgcccagg	300
	gcacgcgcaa	cggggacgcc	accaagggca	tgggcaacga	ggtcaccatc	cacggcatcg	360
10	ctaacgagga	cgccgtccag	ggcatcgcta	acgaggtggc	cgcccagggc	atcgccaacg	420
	aggacgccgc	ccagggaatc	gccgaggatg	tcgcacaggg	catcgccaac	gaggacgccg	480
	cccagggcat	cgccaacaag	gaggccgccc	agggcatcgc	caacgaggac	gccgcccagg	540
	gaatcgctga	ggacgtcgca	cagggcatcg	ccaacgagga	tgccgccag	ggcatcgcca	600
	acgaggaggc	cgcccagggc	atcgccaaca	gggtcgccgc	ccagggcatc	gccaatgacg	660
	ccaccagggg	catcgccgag	gacaccgcca	ggctttnnca	acgacgaacg	ccgtncagg	720
15	cattggttaa	cgaggacgcc	gtcttgggca	ttggccaacg	aacnacgccg	tncaaggcat	780
	tngttaaatg	aaaaaatgga	gttccaccgg	tattcgaaata	accaaggaca	cccgnccaag	840
	ggcattggnc	naactgggga	cttcgcgtcca	agggcctttn	cccaangggg	gacccccgcc	900
	caagggccct	cctttaatgg	gggtcgncgg	nccanngggc	ttnttttacn	ggggaccccc	960
20	tccaangggc	atnttttttt	ttnggggncc	ccccccaag	gggttccctt	tganggggaa	1020
	gtttttccac	gggatttttt	taaaaaggga	ccncttccc	ngggcttttt	ttttanaaag	1080
	gacccatttc	aantttttgn	ttgnaaaggg	accncttcc	ngggtttant	aaanngggac	1140
	ccnccccang	ggntttatta	aattggaanc	ccccccang	gnttttttta	ttnggacccc	1200
	c						1201

<210> 16

<211> 748

<212> DNA

<213> Homo sapiens

<400> 16

	gagtctacgg	cattgctgag	gacgctgccc	agggcatcgc	taatgaggac	gccgaccagg	60
	gcacgcgttaa	tgaggacacc	accagtgca	tcgccaacga	ggaagccgcc	cagggcatcg	120
	ccgaggacgc	catccagggc	atcgccaacg	aggaggttgc	ccagggcatc	gccaatgggg	180
	tcgccgcaca	gggcatcgcc	aatgaggacg	ccaccagggg	catcgccaac	tgggacgccg	240
35	tcacggcctt	cgccaacggg	gacgccgtcc	tcagcttcgc	caacggggac	gccgcccagg	300
	gcacgcgcaa	cggggacgcc	accaagggca	tgggcaacga	ggtcaccatc	cacggcatcg	360
	ctaacgagga	cgccgtccag	ggcatcgcta	acgaggtggc	cgcccagggc	atcgccaacg	420
	aggacgccgc	ccagggaatc	gccgaggatg	tcgcacaggg	catcgccaac	gaggacgccg	480
	cccagggcat	cgccaacaag	gaggccgccc	agggcatcgc	caacgaggac	gccgcccagg	540
40	gaatcgctga	ggacgtcgca	cagggcatcg	ccaacgagga	tgccgccag	ggcatcgcca	600
	acgaggaggc	cgcccagggc	atcgccaaca	gggtcgccgc	ccagggcatc	gccaatgacg	660
	ccaccagggg	catcgccgag	gacaccgcca	ggctttnnca	acgacgaacg	ccgtncagg	720
	cattggttaa	cgaggacgcc	gtcttggg				748

<210> 17

<211> 1232

<212> DNA

<213> Homo sapiens

<400> 17

	ctgaggctgg	ggctggggct	ggggctgagg	ctggagctgg	gactgaggct	ggggctgggg	60
	ctggggctgg	ggctgaggct	ggggctgggg	ctggggctgg	ggctgggact	gaggctgggg	120
	ctggggctga	ggctggggct	gggactgagg	ctggggctgg	gactgaggct	ggggctgggg	180
	ctgagggttg	ggctgggact	gaggctgggg	ctanggtctg	ggctgaggct	ggggctaggg	240
55	ctnaggctga	ggttggggct	ggggctggng	ctgacgtctg	ggctgaggct	nggnetgagg	300
	ctggagctgg	ggctgangct	ggggctgggg	ctgnngctga	ngtggggctg	aggctccngc	360
	tgaagctgag	gctggggcnt	aacgctgagc	tnngngctgg	tgctnatgct	tgnetnanaa	420
	tgngnatgnn	ctnggctnn	cntccnngac	aaananttnn	aacttgnggt	ttnttcctgg	480
	gaatnnaaat	ntccaccann	tntgnaaant	tangcnnttn	ggacnaanaa	anantcnnaa	540
60	antctaanne	cncnanaana	tnctaggana	tgtttacaca	agcaannatn	tnancanac	600
	annccnctc	ntttaaannt	gnattnaaaa	naaanantga	aangnccn	ttnancn	660
	ttnttaantn	gnnaacntna	ctnactnnca	nanatnttaa	aantnggaaa	caancacacn	720
	ntttnanacn	nctnacttcg	gagaataaan	actcnnccctn	nnaatgnctc	agacnaccn	780

```

ntcnttngng cacnnnaaaa tnanancett cttnttttga taccnnnaaa aaanaaaaaac 840
cacttttnaan aannntttta ttcnnaatnn cnannntnta canaggntnt tcacattctn 900
ancnnatttn tccanntnta ttntnccctn ttnnncnnat attnnncana ananantnnn 960
cnnnnnnnacn nncnccnta nnaatattgc acaacnnaan aatannacnn nnttntataa 1020
5 aaatcanaan antancacna cncnnnatc cctanaagtg nttaaaaactc tatgtncnnc 1080
nntctntaat ntannncaaaa tanannnctn nttggnnnat caccannacn tnnnanaccc 1140
nanncttant annntacnn cagcnnnann tncctnnntn tntntnnana acccaactcc 1200
cttatttnat ancanntcac tctccntat cn 1232

```

10 <210> 18  
 <211> 387  
 <212> PRT  
 <213> Homo sapiens

15 <400> 18

Met	Tyr	Ser	Met	Met	Met	Glu	Thr	Asp	Leu	His	Ser	Pro	Gly	Gly	Ala
1				5					10					15	
Gln	Ala	Pro	Thr	Asn	Leu	Ser	Gly	Pro	Ala	Gly	Ala	Gly	Gly	Gly	Gly
			20					25					30		
20 Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Ala	Lys	Ala	Asn	Gln
			35					40					45		
Asp	Arg	Val	Lys	Arg	Pro	Met	Asn	Ala	Phe	Met	Val	Trp	Ser	Arg	Gly
			50			55					60				
Gln	Arg	Arg	Lys	Met	Ala	Gln	Glu	Asn	Pro	Lys	Met	His	Asn	Ser	Glu
				70							75				80
25 Ile	Ser	Lys	Arg	Leu	Gly	Ala	Glu	Trp	Lys	Val	Met	Ser	Glu	Ala	Glu
				85					90					95	
Lys	Arg	Pro	Phe	Ile	Asp	Glu	Ala	Lys	Arg	Leu	Arg	Ala	Leu	His	Met
			100					105					110		
30 Lys	Glu	His	Pro	Asp	Tyr	Lys	Tyr	Arg	Pro	Arg	Arg	Lys	Thr	Lys	Thr
			115				120					125			
Leu	Leu	Lys	Lys	Asp	Lys	Tyr	Ser	Leu	Ala	Gly	Gly	Leu	Leu	Ala	Ala
			130			135					140				
Gly	Ala	Gly	Gly	Gly	Gly	Ala	Ala	Val	Ala	Met	Gly	Val	Gly	Val	Gly
			145			150				155					160
Val	Gly	Ala	Ala	Pro	Val	Gly	Gln	Arg	Leu	Glu	Ser	Pro	Gly	Gly	Ala
				165				170						175	
Ala	Gly	Gly	Ala	Tyr	Ala	His	Val	Asn	Gly	Trp	Ala	Asn	Gly	Ala	Tyr
			180					185					190		
40 Pro	Gly	Ser	Val	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Met	Met	Gln	Glu
			195				200					205			
Ala	Gln	Leu	Ala	Tyr	Gly	Gln	His	Pro	Gly	Ala	Gly	Gly	Ala	His	Pro
			210			215					220				
His	Arg	Thr	Pro	Ala	His	Pro	His	Pro	His	His	Pro	His	Ala	His	Pro
			225			230				235					240
His	Asn	Pro	Gln	Pro	Met	His	Arg	Tyr	Asp	Met	Gly	Ala	Leu	Gln	Tyr
				245					250					255	
Ser	Pro	Ile	Ser	Asn	Ser	Gln	Gly	Tyr	Met	Ser	Ala	Ser	Pro	Ser	Gly
			260					265					270		
50 Tyr	Gly	Gly	Leu	Pro	Tyr	Gly	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	His
			275				280					285			
Gln	Asn	Ser	Ala	Val	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ser	Ser
			290			295					300				
Gly	Ala	Leu	Gly	Ala	Leu	Gly	Ser	Leu	Val	Lys	Ser	Glu	Pro	Ser	Gly
				310						315					320
55 Ser	Pro	Pro	Ala	Pro	Ala	His	Ser	Arg	Ala	Pro	Cys	Pro	Gly	Asp	Leu
				325					330					335	
Arg	Glu	Met	Ile	Ser	Met	Tyr	Leu	Pro	Ala	Gly	Glu	Gly	Gly	Asp	Pro
			340					345					350		
60 Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Gln	Ser	Arg	Leu	His	Ser	Leu	Pro
			355				360					365			
Gln	His	Tyr	Gln	Gly	Ala	Gly	Ala	Gly	Val	Asn	Gly	Thr	Val	Pro	Leu
			370			375					380				

Thr His Ile  
385

5 <210> 19  
<211> 317  
<212> PRT  
<213> Homo sapiens

10 <400> 19  
Met Tyr Asn Met Met Glu Thr Glu Leu Lys Pro Pro Gly Pro Gln Gln  
1 5 10 15  
Thr Ser Gly Gly Gly Gly Asn Ser Thr Ala Ala Ala Ala Gly Gly  
20 25 30  
Asn Gln Lys Asn Ser Pro Asp Arg Val Lys Arg Pro Met Asn Ala Phe  
35 40 45  
15 Met Val Trp Ser Arg Gly Gln Arg Arg Lys Met Ala Gln Glu Asn Pro  
50 55 60  
Lys Met His Asn Ser Glu Ile Ser Lys Arg Leu Gly Ala Glu Trp Lys  
65 70 75 80  
20 Leu Leu Ser Glu Thr Glu Lys Arg Pro Phe Ile Asp Glu Ala Lys Arg  
85 90 95  
Leu Arg Ala Leu His Met Lys Glu His Pro Asp Tyr Lys Tyr Arg Pro  
100 105 110  
25 Arg Arg Lys Thr Lys Thr Leu Met Lys Lys Asp Lys Tyr Thr Leu Pro  
115 120 125  
Gly Gly Leu Leu Ala Pro Gly Gly Asn Ser Met Ala Ser Gly Val Gly  
130 135 140  
Val Gly Ala Gly Leu Gly Ala Gly Val Asn Gln Arg Met Asp Ser Tyr  
145 150 155 160  
30 Ala His Met Asn Gly Trp Ser Asn Gly Ser Tyr Ser Met Met Gln Asp  
165 170 175  
Gln Leu Gly Tyr Pro Gln His Pro Gly Leu Asn Ala His Gly Ala Ala  
180 185 190  
35 Gln Met Gln Pro Met His Arg Tyr Asp Val Ser Ala Leu Gln Tyr Asn  
195 200 205  
Ser Met Thr Ser Ser Gln Thr Tyr Met Asn Gly Ser Pro Thr Tyr Ser  
210 215 220  
Met Ser Tyr Ser Gln Gln Gly Thr Pro Gly Met Ala Leu Gly Ser Met  
225 230 235 240  
40 Gly Ser Val Val Lys Ser Glu Ala Ser Ser Ser Pro Pro Val Val Thr  
245 250 255  
Ser Ser Ser His Ser Arg Ala Pro Cys Gln Ala Gly Asp Leu Arg Asp  
260 265 270  
45 Met Ile Ser Met Tyr Leu Pro Gly Ala Glu Val Pro Glu Pro Ala Ala  
275 280 285  
Pro Ser Arg Leu His Met Ser Gln His Tyr Gln Ser Gly Pro Val Pro  
290 295 300  
Gly Thr Ala Ile Asn Gly Thr Leu Pro Leu Ser His Met  
305 310 315

50 <210> 20  
<211> 443  
<212> PRT  
<213> Homo sapiens

55 <400> 20  
Met Arg Pro Val Arg Glu Asn Ser Ser Gly Ala Arg Ser Pro Arg Val  
1 5 10 15  
Pro Ala Asp Leu Ala Arg Ser Ile Leu Ile Ser Leu Pro Phe Pro Pro  
20 25 30  
60 Asp Ser Leu Ala His Arg Pro Pro Ser Ser Ala Pro Thr Glu Ser Gln  
35 40 45  
Gly Leu Phe Thr Val Ala Ala Pro Ala Pro Gly Ala Pro Ser Pro Pro

[illegible]

		50				55					60						
		Arg 65	Ala	Met	His	Met	Lys 70	Glu	His	Pro	Asp	Tyr 75	Lys	Tyr	Arg	Pro	Arg 80
5		Arg	Lys	Pro	Lys	Thr 85	Leu	Leu	Lys	Lys	Asp 90	Lys	Phe	Ala	Phe	Pro	Val
		Pro	Tyr	Gly	Leu 100	Gly	Gly	Val	Ala	Asp 105	Ala	Glu	His	Pro	Ala	Leu	Lys
		Ala	Gly	Ala	Gly 115	Leu	His	Ala	Gly 120	Ala	Gly	Gly	Gly	Leu 125	Val	Pro	Glu
10		Ser 130	Leu	Leu	Ala	Asn	Pro	Glu 135	Lys	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala
		Ala	Ala	Arg	Val	Phe	Phe	Pro 150	Gln	Ser	Ala	Ala	Ala	Ala	Ala	Ala	Ala
		Ala	Ala	Ala	Ala	Ala	Ala	Gly 165	Ser	Pro	Tyr 170	Ser	Leu	Leu	Asp	Leu	Gly
15		Ser	Lys	Met	Ala 180	Glu	Ile	Ser	Ser	Ser	Ser	Ser	Gly	Leu	Pro	Tyr	Ala
		Ser	Ser	Leu	Gly 195	Tyr	Pro	Thr	Ala 200	Gly	Ala	Gly	Ala	Phe 205	His	Gly	Ala
20		Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Gly	Gly	His	Thr	His
		Ser 225	His	Pro	Ser	Pro	Gly 230	Asn	Pro	Gly	Tyr	Met 235	Ile	Pro	Cys	Asn	Cys 240
		Ser	Ala	Trp	Pro	Ser	Pro	Gly	Leu	Gln	Pro	Pro	Leu	Ala	Tyr	Ile	Leu
25		Leu	Pro	Gly	Met 260	Gly	Lys	Pro	Gln	Leu	Asp 265	Pro	Tyr	Pro	Ala	Ala	Tyr
		Ala	Ala	Ala	Leu 275												
30																	
			<210>		22												
			<211>		533												
			<212>		PRT												
			<213>		Homo sapiens												
35																	
			<400>		22												
		Met 1	Leu	Leu	Asp	Ala 5	Gly	Pro	Gln	Phe	Pro 10	Ala	Ile	Gly	Val	Gly 15	Ser
40		Phe	Ala	Arg	His 20	His	His	His	Ser	Ala 25	Ala	Ala	Ala	Ala	Ala	Ala	Ala
		Ala	Glu	Met	Gln 35	Asp	Arg	Glu	Leu 40	Ser	Leu	Ala	Ala	Ala	Gln	Asn	Gly
		Phe	Val	Asp	Ser	Ala	Ala	Ala 55	His	Met	Gly	Ala	Phe 60	Lys	Leu	Asn	Pro
45		Gly 65	Ala	His	Glu	Leu 70	Ser	Pro	Gly	Gln	Ser	Ser 75	Ala	Phe	Thr	Ser	Gln 80
		Gly	Pro	Gly	Ala 85	Tyr	Pro	Gly	Ser	Ala	Ala 90	Ala	Ala	Ala	Ala	Ala	Ala
		Ala	Leu	Gly	Pro 100	His	Ala	Ala	His 105	Val	Gly	Ser	Tyr	Ser	Gly 110	Pro	Pro
50		Phe	Asn	Ser	Thr 115	Arg	Asp	Phe	Leu 120	Phe	Arg	Ser	Ala	Arg	Leu	Pro	Gly
		Thr	Ser	Ala	Pro	Gly	Gly	Gly 135	Gln	His	Gly	Leu	Phe 140	Gly	Pro	Gly	Ala
55		Gly 145	Gly	Leu	His 150	His	Ala	His	Ser	Asp	Ala 155	Gln	Gly	His	Leu	Leu	Phe 160
		Pro	Gly	Leu	Pro 165	Glu	Gln	His	Gly	Pro	His 170	Gly	Ser	Gln	Asn	Val	Leu
		Asn	Gly	Gln	Met 180	Arg	Leu	Gly	Leu 185	Pro	Gly	Glu	Val	Phe	Gly 190	Arg	Ser
60		Glu	Gln	Tyr	Arg 195	Gln	Val	Ala	Ser 200	Pro	Arg	Thr	Asp	Pro	Tyr	Ser	Ala
		Ala	Gln	Leu	His	Asn	Gln	Tyr	Gly	Pro							

	210		215		220
	Asn Met Ala Ala Ala	Ala Ala His His His	His His His His His		
	225	230	235		240
5	Pro Gly Ala Phe Phe	Arg Tyr Met Arg Gln	Gln Cys Ile Lys Gln	Glu	
		245	250		255
	Leu Ile Cys Lys Trp	Ile Asp Pro Glu Gln	Leu Ser Asn Pro Lys	Lys	
		260	265		270
	Ser Cys Asn Lys Thr	Phe Ser Thr Met His	Glu Leu Val Thr His	Val	
		275	280		285
10	Ser Val Glu His Val	Gly Gly Pro Glu Gln	Ser Asn His Val Cys	Phe	
		290	295		300
	Trp Glu Glu Cys Pro	Arg Glu Gly Lys Pro	Phe Lys Ala Lys Tyr	Lys	
		305	310		315
	Leu Val Asn His Ile	Arg Val His Thr Gly	Glu Lys Pro Phe Pro	Cys	
15		325	330		335
	Pro Phe Pro Gly Cys	Gly Lys Val Phe Ala	Arg Ser Glu Asn Leu	Lys	
		340	345		350
	Ile His Lys Arg Thr	His Thr Gly Glu Lys	Pro Phe Gln Cys Glu	Phe	
		355	360		365
20	Glu Gly Cys Asp Arg	Arg Phe Ala Asn Ser	Ser Asp Arg Lys Lys	His	
		370	375		380
	Met His Val His Thr	Ser Asp Lys Pro Tyr	Leu Cys Lys Met Cys	Asp	
		385	390		395
	Lys Ser Tyr Thr His	Pro Ser Ser Leu Arg	Lys His Met Lys Val	His	
25		405	410		415
	Glu Ser Ser Pro Gln	Gly Ser Glu Ser Ser	Pro Ala Ala Ser Ser	Gly	
		420	425		430
	Tyr Glu Ser Ser Thr	Pro Pro Gly Leu Val	Ser Pro Ser Ala Glu	Pro	
		435	440		445
30	Gln Ser Ser Ser Asn	Leu Ser Pro Ala Ala	Ala Ala Ala Ala Ala		
		450	455		460
	Ala Ala Ala Ala Ala	Ala Ala Val Ser Ala	Val His Arg Gly Gly	Gly	
		465	470		475
	Ser Gly Ser Gly Gly	Ala Gly Gly Gly Ser	Gly Gly Gly Ser Gly	Ser	
35		485	490		495
	Gly Gly Gly Gly Gly	Gly Ala Gly Gly Gly	Gly Gly Gly Ser Ser	Gly	
		500	505		510
	Gly Gly Ser Gly Thr	Ala Gly Gly His Ser	Gly Leu Ser Ser Asn	Phe	
		515	520		525
40	Asn Glu Trp Tyr Val				
		530			